

26


MARCH 2017, 8 (1): 45-312

www.biodiversityjournal.com

ISSN 2039-0394 (Print Edition)

ISSN 2039-0408 (Online Edition)

with the support of



Biodiversity Journal

MONOGRAPH

PROCEEDINGS OF THE 3rd INTERNATIONAL
CONGRESS

“BIODIVERSITY, MEDITERRANEAN,
SOCIETY”

SEPTEMBER 4th-6th, 2015
NOTO-VENDICARI, ITALY



Juniperus turbinata Guss. - Piano Pirrera, Acate, Sicily

***Juniperus turbinata* Guss. (Cupressaceae).** *Juniperus phoenicea* was described by Linnaeus (1753) from specimens collected from Montpellier, southern France; later Gussone (1845), described *J. turbinata* from southern Sicily differentiating it from the previous one. After many taxonomic vicissitudes, the two species are currently recognized by various authors and by the IUCN's Conifer Specialist Group. *Juniperus turbinata* differs from *J. phoenicea* mostly by having a larger (12-14 mm), sub-ovoid cone and leaves without scarious border (Arrigoni, 2012). *J. phoenicea* is a western European taxon that grows in a wide bioclimatic range from thermo- to supramediterranean belt with semiarid to humid ombrotype, in Spain, southern France, Italy in Maritime and Apuane Alps, while *J. turbinata* is found from the infra- to supramediterranean belt with arid to humid ombrotype and is widespread in the Mediterranean region and Canary Islands, both in coastal and inland regions (Mazur et al., 2016). *J. turbinata* is a large shrub or a small tree (8-12m tall), which grows mainly in coastal areas of the Mediterranean region and in mountains of northwest of Africa. In Sicily, the species is generally rare but a large population was recently discovered near Acate in SE Sicily (Minissale & Sciandrello, 2013). It is a typical Mediterranean species globally evaluated as Near Threatened (NT) by the IUCN world red list (Farjon, 2013). In fact although it is widely distributed along the coasts of the Mediterranean, subpopulations are scattered and usually small or very small. It often occurs close to urbanized coastal strips, or paleo-dunes where much of the original habitat has been destroyed. So reinforcement of *J. turbinata* populations and habitat restoration should be implemented.

References: Arrigoni PV., 2012. Miscellaneous notes about some taxa of the Italian flora. *Webbia*, 67: 3746. - Farjon A., 2013. *Juniperus turbinata*. The IUCN Red List of Threatened Species 2013: e.T16349692A84434669. Downloaded on 26 March 2017. - Gussone J., 1845. *Florae Siculae Synopsis* 2, Neapoli, p. 634 - Linnaeus C., 1753. *Species plantarum* 2. Stockholm: Laurentii Salvii, p. 640. - Mazur M., Minissale P., Sciandrello S. & Boratyński A., 2016. Morphological and ecological comparison of populations of *Juniperus turbinata* Guss. and *J. phoenicea* L. from the Mediterranean region. *Plant Biosystems*, 150: 313-322. - Minissale P. & Sciandrello S., 2013. A relic wood of *Juniperus turbinata* Guss. (Cupressaceae) in Sicily: Structural and ecological features, conservation perspectives. *Plant Biosystems*, 147: 145-157.

Pietro Minissale, Dipartimento di Scienze Biologiche, Geologiche e Scienze Ambientali, Università di Catania, Via A. Longo 19, 95125 Catania, Italy; e-mail: p.minissale@unict.it - Cover photo by Pietro Minissale