

# Metallothioneins as a biomarker of metallic pollution in *Donax trunculus* Linnaeus, 1758 (Mollusca Bivalvia) from the Gulf of Annaba (Algeria)

Amina Yezli, Abdelmalek Salahi, Asma Boukari & Nouredine Soltani\*

Laboratory of Applied Animal Biology, Badji Mokhtar University, 23000 Annaba, Algeria

\*Corresponding author: [nouredinesoltani@univ-annaba.org](mailto:nouredinesoltani@univ-annaba.org); [nouredinesoltani36@gmail.com](mailto:nouredinesoltani36@gmail.com)

## ABSTRACT

In recent years, the Gulf of Annaba has observed an increase in the pollution level, in particular by heavy metals. The present experiment aimed to assess pollution by measuring the rate of metallothioneins (MTs), a biomarker of metallic contamination, measured in the digestive gland of *Donax trunculus* Linnaeus, 1758 (Mollusca Bivalvia) an edible mollusk widely used in biomonitoring of water quality. This sentinel species was sampled during two seasons (winter and summer 2019) at two sites in the Gulf of Annaba (Northeast Algeria): El Battah site far from any source of pollution and Sidi Salem site exposed to various sources of pollution. The biochemical analyzes reveal spatio-temporal changes. The levels of proteins recorded in the two studied sites were higher in summer and values from El Battah individuals were significantly higher compared to those from Sidi Salem. In addition, they show an induction in MT rates in individuals from the two sites (Sidi Salem and El Battah) in summer with higher values recorded in individuals from Sidi Salem. This difference is related to the proximity of sources of contamination at Sidi Salem compared to El Battah. The rates of MTs vary also according to abiotic factors such as hydrodynamics or temperature which can explain the significant seasonal difference with a more marked induction in summer. Conclusively, our study shows the sensitivity of *D. trunculus* to chemical stress induced by heavy metal contamination as evidenced by an induction of MTs with significant effects of both sites and seasons.

## KEY WORDS

Pollution; Gulf of Annaba; *Donax trunculus*; Biomarker; Proteins; Metallothioneins.

Received 31.05.2022; accepted 24.09.2022; published online 24.10.2022

## INTRODUCTION

The development of human activities is responsible for the chronic pollution of the environment by organic and inorganic contaminants (Larba & Soltani, 2014; Breitwieser et al., 2020; Douafer et al., 2020). The marine environment, and in particular coastal areas, receives a large amount of contaminants from urban, agricultural, port and industrial activities. These contaminants

cause increased pollution of marine ecosystems and have a significant toxic impact on the health of living organisms (Amamra et al., 2019; Hamdani et al., 2020). However, the increase in ocean pollution caused by toxic metals has resulted in bioaccumulation of toxic metals in marine animals, particularly bivalves such as clams, and the consequent toxicity associated with heavy metals in these animals (Mengxu et al., 2020). Thus, biomonitoring is an important tool to determine the

link between current pollution levels and observed effects in the field (Asker et al., 2016). Bivalves belong to the first-choice species as bioindicators for environmental and chemical stress. They are sentinel benthic organisms living as filter feeders and exposed to different environmental compartments (Laitano et al., 2016; Dos Santos et al., 2022).

Biomarkers are an important tool for understanding the possible harmful effects of pollutants on organisms (Boshoff et al., 2015). They are inhibition biomarkers such as metallothioneins (MTs) which are widely used as a biomarker of metal contamination by binding and removing toxic metals (Pedrini-Martha et al., 2017). A defense biomarker like MTs was considered in this study to investigate metal accumulation. MTs are low molecular weight, cysteine-rich proteins induced in response to various environmental stressors, including metal ions, cytokines and free radicals (Coyle et al., 2002). MTs are involved in metal uptake, storage and excretion (Breitwieser et al., 2020). Their concentrations in whole soft tissues or in a particular bivalve tissue have been widely used in ecotoxicological studies and biomonitoring programs (Yen Lea et al., 2016; Merad & Soltani, 2017; Pedrini-Martha et al., 2017).

In this context, the main objective of this work was to study the in situ seasonal responses (winter and summer) of a biochemical biomarker MTs in *D. trunculus*, a locally prevalent edible mollusk sampled at two sites in the Gulf of Annaba: a pol-

luted site (Sidi Salem) and a relatively clean site (El Battah).

## MATERIAL AND METHODS

### Study area

The Gulf of Annaba is located in the Northeast of Algeria. It is limited by the Rosa Cap ( $8^{\circ}14'13''$  E,  $36^{\circ}56'46''$  N) to the east and by the Gard Cap ( $7^{\circ}16'00''$  E,  $36^{\circ}58'02''$  N) to the west.

The El Battah site ( $36^{\circ}5'49''$  N,  $7^{\circ}56'57''$  E), is located about 30 km to the east of Annaba far from any human activities and is considered as a relatively clean site because of its remoteness from anthropogenic activity, its important hydrodynamic exposure, and the species' abundance (Rabei et al., 2018; Boukari et al., 2021). Sidi Salem site ( $36^{\circ}51'19''$  N,  $7^{\circ}47'48''$  E), located about 1 km to the east of Annaba city, receives industrial and domestic wastewater, and is considered as the polluted area (Fig. 1).

### Samples collection

Mollusk bivalves (*D. trunculus*) with the same shell length ( $25 \pm 1$  mm) (Hamdani et al., 2020) were collected seasonally in the winter and summer of 2019, from two sampling sites (El Battah and Sidi Salem). Animals were transported in cold boxes to the laboratory: the dissection and the sam-



Figure 1. Location of sampling sites of *D. trunculus* in the Gulf of Annaba: the reference site (El Battah) and the polluted site (Sidi Salem).

pling of the digestive glands are carried out the same day of the fishing.

**Physico-chemical parameters**

The physico-chemical parameters of seawater were recorded using a multiparameter (Multi 340 i, Germany) during the study period at the two sites, El Battah and Sidi Salem. The following parameters were considered: temperature (°C), pH, salinity (PSU) and dissolved oxygen (%) (Table 1).

**Protein quantification**

The proteins were extracted following the procedure of Shibko et al. (1966) recently described (Boukari et al., 2021) and quantified according to the method of Bradford (1976). This method consists in measuring protein concentrations in the supernatants by using bovine serum albumin (Sigma) as a standard and Coomassie Brilliant Blue (BBC) as a reagent. The absorbance was read at a wavelength of 595 nm. The assay was conducted with 5 repeats.

**Determination of metallothionein rate**

The determination of metallothioneins in digestive glands was performed following the procedure of Viarengo et al. (1997) previously described (Amira et al., 2018). This method is based on the quantification of SH-residue contents by a spectrophotometric method using Ellman’s reagent. Clams were collected (5 specimens from each site). Each digestive gland was dissected and analyzed individually. Absorbance was read at 412 nm and MT levels quantified by use glutathione (Sigma) as a standard. The results are expressed as µg/mg proteins.

**Statistical Analysis**

Data are represented as mean ± standard deviation (SD). The statistical analyses were performed using prism version 7 for Windows (GraphPad software, La Jolla California, USA, www.Graphpad.com). The comparison of the mean values between sites was made by Student’s t-test. The combined effects of seasons and sites were investigated using two-way analysis of variance (ANOVA) followed by a post-hoc HSD Tukey test.

**RESULTS**

The physico-chemical parameters of the sea water show average temperature values of 15.12±0.60 °C at El Battah and 14.54±0.31 °C at Sidi Salem. The value of pH is 7.83±0.20 at the site of El Battah and 7.87±0.12 at the site Sidi Salem. Concerning salinity a value of 22.50±0.45 PSU was recorded at the site El Battah and 24.20±0.60 PSU at Sidi Salem. The dissolved oxygen shows a value of 5.51±0.31 % for the site of El Battah and for the site of Sidi Salem 5.37±0.27 %. Statistical analysis by Student’s t-test shows no significant difference between the two studied sites (p>0.05) for all parameters considered (Table 1).

**Protein amounts**

The results of the MTs assay obtained are expressed in terms of a quantity of protein. For this purpose, a regression line expressing the absorbance as a function of the quantity of the standard, albumin, was determined (Fig. 2).

Site Parameters	El Battah	Sidi Salem
Temperature (°C)	15.12±0.60 a	14.54±0.31 a
pH	7.83±0.20 a	7.87±0.12 a
Salinity (PSU)	22.50±0.45 a	24.20±0.60 a
Dissolved oxygen (%)	5.51±0.31 a	5.37±0.27 a

Table 1. Physic-chemical parameters of water at the two sampling sites. For each parameter, mean values followed by the same letter are not significantly different (p> 0.05).

Site Season	El Battah	Sidi Salem
Winter	79.61 ± 0.90 a A	71.68 ± 1.60 b A
Summer	96.10 ± 1.7 a B	80.72 ± 0.70 b B

Table 2. Total protein amounts (µg/mg tissue) in the digestive gland of *D. trunculus* during winter and summer 2019 (m ± SD; n= 5; For the same season, means followed by the same lowercase letter are not significantly different, while for the same site, means followed by the same uppercase letter are not significantly different at p > 0.05).

Biochemical analyses reveal a significant ( $p < 0.001$ ) increase in proteins levels at Sidi Salem compared to El Battah during the two seasons (Table 2). Maximum values were recorded in El Battah site with an average of  $96.10 \pm 1.7$  ( $\mu\text{g}/\text{mg}$  tissue) in summer against minimum values in Sidi Salem site with an average rate  $80.72 \pm 0.70$  ( $\mu\text{g}/\text{mg}$  tissue). A two-way ANOVA revealed significant ( $p < 0.001$ ) effects site ( $F_{1, 12} = 14.92$ ;  $P < 0.0023$ ), season ( $F_{(1, 12)} = 12.42$ ;  $P < 0.0042$ ) and interaction site season was no significant ( $F_{1, 12} = 1.271$ ;  $P < 0.2817$ ).

### Metallothionein rates

The results obtained concerning the levels of




Figure 2. Protein assay: regression line expressing absorbance as a function of the amount of albumin ( $\mu\text{g}$ ) ( $R^2$  coefficient of determination).

Site Season	El Battah	Sidi Salem
Winter	$2.19 \pm 0.53$ a A	$2.69 \pm 0.37$ a A
Summer	$3.03 \pm 0.38$ b B	$4.10 \pm 0.67$ b B

Table 3. Metallothionein rates ( $\mu\text{g}/\text{mg}$  proteins) in the digestive gland of *D. trunculus* during winter and summer 2019 ( $m \pm \text{SD}$ ;  $n = 5$ ; For the same season, means followed by the same lowercase letter are not significantly different, while for the same site, means followed by the same uppercase letter are not significantly different at  $p > 0.05$ ).

protein. For this purpose, a regression line expressing the absorbance as a function of the quantity of the standard, albumin, was determined (Fig. 2). MTs are summarized in Table 3. They reveal a significant increase ( $p < 0.001$ ) in the levels of MTs observed in Sidi Salem compared to El Battah during both two seasons. The maximum values were recorded at the Sidi Salem site with a mean of  $4.10 \pm 0.67$  ( $\mu\text{g}/\text{mg}$  proteins) in summer, compared to the minimum values at the El Battah site with a mean level of  $3.03 \pm 0.38$  ( $\mu\text{g}/\text{mg}$  protein). A two-way ANOVA revealed significant ( $p < 0.001$ ) effect of site ( $F_{1, 16} = 24.47$ ;  $P < 0.0001$ ) and season ( $F_{1, 16} = 12.11$ ;  $P < 0.0031$ ). The site-season interaction was not significant ( $F_{1, 16} = 1.527$ ;  $P < 0.2344$ ).

## DISCUSSION

### Changes in protein amounts

Proteins play an important role in repairing and protecting organisms from environmental damage (Umminger, 1977; Ansell et al., 1980; Maharajan et al., 2012). Protein production and expression reflect physiological responses to changing or stressful environmental conditions (Gotelli et al., 2012; Pedrosa et al., 2017). Under stressful conditions, proteins provide energy in metabolic pathways and biochemical reactions to counter this aggression (Maharajan et al., 2012; Do Nascimento et al., 2016).

Our study conducted in *D. trunculus* shows significant decreases in protein levels over the two seasons in the polluted site Sidi Salem compared to El Battah site. Similar results were reported by Yusseppone et al. (2020) where protein analysis in the digestive gland of *Diplodon chilensis* Gray 1828 (Bivalvia Hyriidae) transplanted into a polluted site showed a significant decrease. Since bivalves have a limited amount of carbohydrates and lipids, the next alternative source of energy to meet the increased demand is protein (Le Gal et al., 1997; Xuan et al., 2011; Yeung et al., 2017). Many bivalve species have shown a reduction in proteins to overcome Cd-induced stress: *Mytilopsis. sallei* (Recluz, 1849) (Bivalvia Dreissenidae) exposed to the concentration of 0.71 mg/L (Uma-Devi et al., 1996), *Anodonta anatina* Linnaeus, 1758 (Bivalvia Unionidae) with algae containing the dose of

0.0059 µg/g dry weight (Ngo et al., 2012) or *Mytilus edulis* Linnaeus, 1758 (Bivalvia Mytilidae) treated at the concentration of 0.2 mg/L (Géret et al., 2002). The depletion of total protein levels suggests increased proteolysis and possible use of their degradation products for metabolic purposes, and may be due to increased catabolism and decreased anabolism of proteins, as reported in studies on the freshwater bivalve *Parreysia corrugata* (O.F. Müller, 1774) (Bivalvia Unionidae) (Deshmukh & Lomte, 1998).

### Changes in metallothionein rates

Metallothionein levels were determined in the digestive gland of *D. trunculus* in individuals sampled at the El Battah site as well as at Sidi Salem in winter and summer of 2019; our data showed highest values recorded in summer. The statistical analysis reveals that there are significant effects of season and site, confirming previous observations made by Rabei et al. (2018) and Amira et al. (2018) in the same species at the same sites in the Gulf of Annaba. This response is related to the higher metal pollution levels in Sidi Salem compared to El Battah (Belabed et al., 2017; Amira et al., 2018; Rabei et al., 2018; Drif et al., 2019). In fact, the Gulf of Annaba has a strong industrial activity which contributes to the discharge of multiple pollutants such as heavy metals (Abdenour et al., 2000; Beldi et al., 2006; Larba & Soltani, 2014; Belabed et al., 2017, Amira et al., 2018; Drif et al., 2019) and the site of Sidi Salem is exposed to various pollutants from domestic, agricultural and industrial sources (Ounissi et al., 2016; Rabei et al., 2018; Kheroufi et al., 2021, Kroini et al., 2021). The relationship between the concentrations of heavy metals and MTs depends on the concentration and chemical speciation of the accumulated metals (Amira et al., 2018).

A recent report conducted in the same species also showed that MTs are induced by Cd with a concentration effect in several tissues (adductor muscle, gill, digestive gland) after 96 hours of exposure to various concentrations (Moncaleano-Niño et al., 2017). In the same biological model, Tlili et al. (2013) noted an increase in digestive gland MT levels of *D. trunculus* at a site close to the chemical and metallurgical industries in Tunisia. Sand grain size also affects metal availability (Amira et al., 2018; Manju et al., 2020). This explains the in-

crease in rates at the Sidi Salem site, which is made up of fine sand, in contrast to the El Battah site, which is made up of medium sand (Amira et al., 2018). A similar induction has been recorded in Morocco where the proximity of the Mohammedia site (Morocco) to the discharge of a petrochemical company using metals (mainly mercury) among mussels *Mytilus galloprovincialis* (El Haimeur et al., 2017). The study by Mdelgi-Lasram et al. (2007) on clams *Ruditapes decussatus* Linnaeus, 1758 (Bivalvia Veneridae) from the Bizerte lagoon (Tunis) at three sites shows that the levels of MTs in the digestive gland are different at the three sites, with a positive correlation with the degree of metal pollution of sites.

Moreover, the comparison of MTs rates between the two seasons (winter and summer) at the same sites (Sidi Salem and El Battah) reveals an increase in MTs rates in summer compared to winter for both sites. This difference may be linked to the fact that in winter, sea currents are stronger than in summer, thus contributing to the dilution of pollutants (Khelifi et al., 2007; Ounissi et al., 2016). However, a change in abiotic factors can affect the bioavailability of pollutants to living organisms and their penetration into tissues (Nahrgang et al., 2013). Indeed, increasing water temperature leads to an increase in the solubility of heavy metals (Hertika et al., 2018). Similarly, increasing salinity will have the same impact on the concentration of these elements (Kavun et al., 2002) and the decrease in dissolved oxygen concentration increases their toxicity (Rotchell et al., 2001). Furthermore, this difference may be related to increased wastewater and air emissions (Qaderi et al., 2018). In addition, other authors have suggested that MT levels may increase during the reproductive period and throughout cell proliferation and differentiation (Coyle et al., 2002; Boulajfene et al., 2019; Hamdani et al., 2020).

### CONCLUSIONS

This paper reports the results of a biochemical biomarker MTs measured in the digestive gland of *D. trunculus*, a useful sentinel for environmental monitoring. The results will contribute to the development of an environmental monitoring program based on relevant bioindicators. The level of metal-

lic pollution was found relatively high at Sidi Salem compared to El Battah.

## ACKNOWLEDGEMENTS

This work was supported by the National Fund for Scientific Research to Pr. N. Soltani (Laboratory of Applied Animal Biology) and the Ministry of High Education and Scientific Research of Algeria (PRFU Project N° D01N01UN230120190009 to Pr. N. Soltani).

## REFERENCES

- Abdenmour C., Smith B.D., Boulakoud M.S., Samraoui B. & Rainbow P.S., 2000. Trace metals in shrimps and sediments from Algerian water. *Journal of Catalysis Material Environmental*, 3: 9–12.
- Amamra F., Sifi K., Kouachi N. & Soltani N., 2019. Evaluation of the impact of pollution in the gulf of Annaba (Algeria) by measurement of environmental stress biomarkers in an edible mollusk bivalve *Donax trunculus*. *Fresenius Environmental Bulletin*, 28: 908–915.
- Amira A., Merad I., Almeida C.M.R., Guimara L. & Soltani N., 2018. Seasonal variation in biomarker responses of *Donax trunculus* from the Gulf of Annaba (Algeria): Implication of metal accumulation in sediments. *Comptes Rendus Geoscience*, 350:173–179.
- Ansell A.D., Frenkiel L. & Mouëza M., 1980. Seasonal changes in tissue weight and biochemical composition for the bivalve *Donax trunculus* L. on the Algerian Coast. *Journal of Experimental Marine Biology and Ecology*, 45: 105–116.
- Asker N., Albertsson E., Wijkmark E., Bergek S., Parkkonen J., Kammann U., Holmqvist I., Kristiansson E., Strand J., Gercken J. & Forlin L., 2016. Biomarker responses in eelpouts from four coastal areas in Sweden, Denmark and Germany. *Marine Environmental Research*, 120: 32–43.
- Belabed B., Meddour A., Samraoui B. & Chenchouni H., 2017. Modeling seasonal and spatial contamination of surface waters and upper sediments with trace metal elements across industrialized urban areas of the Seybouse watershed in North Africa. *Environmental Monitoring and Assessment*, 189: 1–19.
- Beldi H., Gimbert F., Maas, S., Scheifler R. & Soltani N., 2006. Seasonal variations of Cd, Cu, Pb and Zn in the edible mollusc *Donax trunculus* (Mollusca, Bivalvia) from the Gulf of Annaba, Algeria. *African Journal of Agricultural Research*, 1: 85–90.
- Boshoff M., Jordaens K., Baguet S. & Bervoets L., 2015. Trace metal transfer in a soil–plant–snail microcosm field experiment and biomarker responses in snails. *Ecological Indicators*, 48: 636–648.
- Boukari A., Hamoudi F.S. & Soltani N., 2021. Biochemical modification in an edible Mollusk (*Donax trunculus*) during transplantation into polluted environment. *Fresenius Environmental Bulletin*, 30: 2416–2422.
- Boulajfene W., Stroglyoudi E., Lasram M., El Mlayah A., Vassiliki-Angelique C. & Zouari-Tlig S., 2019. Biological and biochemical assessment in *Phorcus articulatus* (Lamarck, 1822): contamination and seasonal effect. *Environmental Monitoring and Assessment*, 191(555): 16.
- Bradford M.M., 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the Principle of protein-dye binding. *Anal. Biochem*, 72 : 254–278.
- Breitwieser M., Bruneau M., Barbarin M., Churlaud C., Mouneyrac C. & Thomas H., 2020. Is metallothionein in *Mimachlamys varia* a suitable biomarker of trace elements in the waters of the French Atlantic coast? *Environnemental Science and Pollution Research*: 27: 20259–20272. <https://doi.org/10.1007/s11356-020-08392-1>
- Coyle P. & Philcox J.C. & Rofe A.M., 2002. Metallothionein: The multipurpose protein. *Cellular and Molecular Life Sciences*, 59: 627–647.
- Deshmukh R. & Lomte, A., 1998. Effect of heavy metal (CuSO<sub>4</sub>) on protein activity of freshwater bivalve, *Parreysia corrugata*. *Journal of Ecotoxicology Monitoring*, 16: 704–708.
- Do Nascimento L.F., da Silveira L.C., Nisembaum L.G., Colquhoun A., Abe AS., Mandarim-de-Lacerda C.A. & de Souza S.C., 2016. Morphological and metabolic adjustments in the small intestine to energy demands of growth, storage, and fasting in the first annual cycle of a hibernating lizard (*Tupinambis merianae*). *Comparative Biochemistry and Physiology, Part A: Molecular and Integrative Physiology*, 195: 55–64.
- Dos Santos F. Silva., Neves R.A.F., Crapez M.A.C., Teixeira V.L. & Krepsky N., 2022. How does the brown mussel *Perna perna* respond to environmental pollution? A review on pollution biomarkers. *Journal of Environmental Sciences*, 111: 412–428.
- Douafer L., Zaidi N. & Soltani N., 2020. Seasonal variation of biomarker responses in *Cantareus aspersus* and physico-chemical properties of soils from Northeast Algeria. *Environmental Science and Pollution Research*, 27: 24145–24161. <https://doi.org/10.1007/s11356-020-08694-4>.
- Drif F., Abdenmour C., Çiğerci I.H. & Muddassir Ali M., 2019. Preliminary Assessment of Stress and Genotoxicity Biomarkers in Bivalve Molluscs from the

- Gulf of Annaba, Algeria. Bulletin of Environmental Contamination and Toxicology, 102: 1–3.
- El Haimour B., Zbiry M., Bouhallaoui M., Elkhiaati . & Benhra A., 2017. Multimarker approach to assess seasonal variation of pollutant effects in *Mytilus galloprovincialis* in the Casablanca region (Moroccan Atlantic coast). Bulletin de le Société zoologique de France, 142: 203–223.
- Géret F., Jouan A., Turpin V., Bebianno M.J. & Cosson R.P., 2002. Influence of metal exposure on metallothionein synthesis and lipid peroxidation in two bivalve mollusks: the oyster (*Crassostrea gigas*) and the mussel (*Mytilus edulis*). Aquatic Living Resources, 15: 61–66.
- Gotelli N.J., Ellison A.M. & Ballif A., 2012. Environmental proteomics, biodiversity statistics and food-web structure. Trends in Ecology & Evolution, 1522: 1–7.
- Hamdani A., Soltani N. & Zaidi N., 2020. Growth and reproduction of *Donax trunculus* from the Gulf of Annaba (Northeast Algeria) in relation to environmental conditions. Environmental Science and Pollution Research, 27: 41656–41667.
- Hertika AMS., Kusriani K., Indrayani E., Nurdiani R. & Putra RBDS., 2018. Relationship between levels of the heavy metals lead, cadmium and mercury, and metallothionein in the gills and stomach of *Crassostrea iredalei* and *Crassostrea glomerata*. F1000 Research, 7: 1239.  
<http://dx.doi.org/10.1016/j.jes.2021.04.006>.
- Kavun VY., Shulkin VM. & Khristoforova NK., 2002. Metal accumulation in mussels of the Kuril Islands, north-west Pacific Ocean. Marine Environmental Research, 53: 219–226.
- Khélifi-Touhami M., Ounissi M., Saker I., Haridi A., Djorfi S. & Abdenour C., 2007. The hydrology of the Mafrag estuary (Algeria): Transport of inorganic nitrogen and phosphorus to the adjacent coast. Journal of Food Agriculture and Environment, 4: 340–346.
- Kheroufi N., Hamdani A. & Soltani N., 2021. Acute exposure of cadmium on *Donax trunculus* Linnaeus, 1758 (Mollusca Bivalvia) during the vitellogenesis process: histological and biochemical aspects. Biodiversity Journal, 12: 865–873.  
<https://doi.org/10.31396/Biodiv.Jour.2021.12.4.865.873>
- Kroini H., Hamdani A., Soltani N., Zaidi N. & Sleimi N., 2021. Effects of cadmium exposure on proximate analysis and metallic element contents of wedge clam *Donax trunculus*). Applied Ecology and Environmental Research, 19: 5103–5128.  
[http://dx.doi.org/10.15666/aer/1906\\_51035128](http://dx.doi.org/10.15666/aer/1906_51035128)
- Laitano M. & Fernandez-Gimenez A., 2016. Are Mussels Always the Best Bioindicators Comparative Study on Biochemical Responses of Three Marine Invertebrate Species to Chronic Port Pollution? Bulletin of Environmental Contamination and Toxicology, 97: 50–55.
- Larba R. & Soltani N., 2014. Use of the land snail *Helix aspersa* for monitoring heavy metal soil contamination in Northeast Algeria. Environmental Monitoring and Assessment, 186: 4987–4995.
- Le Gal Y.L., Lagadic L., Le Bras S. & Caquet T., 1997. Charge énergétique en adényles (CEA) et autres biomarqueurs associés au métabolisme énergétique. In: Lagadic L., Caquet T., Amiard J.C. & Ramade F., Biomarqueurs en écotoxicologie: aspects fondamentaux. Masson, Paris, pp. 241–285.
- Manju M.N., Ratheesh Kumar C.S., Resmic P., Gireeshkumar T.R., Manju M.J., Salasa P.M. & Chandramohanakumar N., 2020. Trace metal distribution in the sediment cores of mangrove ecosystems along northern Kerala coast, south-west coast of India. Marine Pollution Bulletin, 153: 110946.
- Mdelgi-Lasram E., Fassatoui C., Moraga D. & Romdhane M.S., 2007. Impact saisonnier de l’environnement sur la structure génétique de trois populations de *Ruditapes decussatus* dans l’écosystème lagunaire de bizerte (tunisie). Bulletin de l’Institut des Sciences et Technologie de la Mer de Salammbô, 34: 109–118.
- Mengxu C., Jiayi Z., Jinhuang L., Hongchao T., Yifei S., Alan K.C. & Xueping Y., 2020. Changes in oxidative stress biomarkers in *Sinonovacula constricta* in response to toxic metal accumulation during growth in an aquaculture farm. Chemosphere, 248: 125974.
- Merad I. & Soltani, N., 2017. Sublethal effects of cadmium on energy reserves in the edible Mollusk *Donax trunculus*. Journal of Entomology and Zoology Studies, 5: 100–105.
- Moncaleano-Niño A.M., Barrios-Latorre S.A., Poloche-Hernández J.F., Becquet V., Huet V., Villamil L., Thomas-Guyon H., Ahrens M.J. & Luna-Acosta A., 2017. Alterations of tissue metallothionein and vitellogenin concentrations in tropical cup oysters (*Saccostrea* sp.) following short-term (96h) exposure to cadmium. Aquatic Toxicology, 185: 160–170.
- Nahrgang J., Brooks S.J., Evensen A., Camus L., Jonsson M., Smith T.J., Lukina J., Frantzen M., Giarratano E. & Renau P.E., 2013. Seasonal variation in biomarkers in blue mussel (*Mytilus edulis*), Icelandic scallop (*Chlamys islandica*) and Atlantic cod (*Gadus morhua*) Implications for environmental monitoring in the Barents Sea. Aquatic Toxicology, 127: 21–35.
- Ngo H.T.T., Gerstmann S. & Frank H., 2012. Subchronic effects of environment like cadmium levels on the bivalve *Anodonta anatina* (Linnaeus, 1758): II. Effects on energy reserves in relation to calcium metabolism. Environmental Toxicology and Chemistry, 93: 1802–1814.
- Ounissi M., Laskri H. & Khelifi-Touhami M., 2016. Net-zooplankton abundance and biomass from Annaba

- bay (SW Mediterranean sea) under estuarine influences. *Mediterranean Marine Science*, 17: 519–532.
- Pedrini-Martha V., Schnegg, R., Baurand, P.E., deVaufleury, A. & Dallinger R., 2017. The physiological role and toxicological significance of the non-metal-selective cadmium/coppermetallothionein isoform differ between embryonic and adult helix snails. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 199: 38–47.
- Pedrosa J., Gravato C., Campos D., Cardoso P., Figueira E., Nowak C., Soares A.M.V.M., Barata C. & Pestana J.L.T., 2017. Investigating heritability of cadmium tolerance in *Chironomus riparius* natural populations: A physiological approach. *Chemosphere*, 170: 83–94.
- Qaderi F., Sayahzadeh A.H. & Azizi M., 2018. Efficiency optimization of petroleum wastewater treatment by using of serial moving bed biofilm reactors. *Journal of Cleaner Production*, 192: 665–677.
- Rabei A., Hichami A., Beldi H., Bellenger S., Khan A.N. & Soltani N., 2018. Fatty acid composition, enzyme activities and metallothioneins in *Donax trunculus* (Mollusca, Bivalvia) from polluted and reference sites in the Gulf of Annaba (Algeria): Pattern of recovery during transplantation. *Environmental Pollution*, 237: 900–907.
- Rotchell J.M., Clarke K.R. & Newton L.C., 2001. Hepatic metallothionein as a biomarker for metal contamination: age effects and seasonal variation in European flounders (*Pleuronectes flesus*) from the Severn Estuary and Bristol Channel. *Marine Environmental Research*, 52: 151–171.
- Shibko S., Koivistoinen P., Tratnyek C.A., Newhall A.R. & Friedman L., 1966. A method for sequential quantitative separation and determination of protein, RNA, DNA, lipid, and glycogen from a single rat liver homogenate or from a subcellular fraction. *Analytical Biochemistry*, 19: 514–528.
- Tlili S., Minguez, L., Giamberini L., Geffard A., Boussetta H. & Mouneyrac C., 2013. Assessment of the health status of *Donax trunculus* from the Gulf of Tunis using integrative biomarker indices. *Ecological Indicators*, 32: 285–293.
- Umminger B.L., 1977. Relation of whole blood sugar concentration in vertebrate to standard metabolic rate. *Comparative Biochemistry and Physiology Part A*, 56: 457–460.
- Uma Devi. V., 1996. Bioaccumulation and metabolic effects of cadmium on marine fouling bryozoan bivalve, *Mytilopsis sallei* (Recluz). *Archives of Environmental Contamination and Toxicology*, 31: 47–53.
- Viarengo A., Ponzano E., Dondero F. & Fabbri R., 1997. A simple spectrophotometric method for metallothionein evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. *Marine Environmental Research*, 44: 69–84.
- Xuan R., Wang L., Sun M., Ren G. & Jiang M., 2011. Effects of cadmium on carbohydrate and protein metabolisms in the freshwater crab *Sinopotamon yangtsekiense*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 154: 268–74.
- Yen Lea T.T., Zimmermann S. & Sures B., 2016. How does the metallothionein induction in bivalves meet the criteria for biomarkers of exposure? *Environmental Pollution*, 212: 257–268.
- Yeung J.W.Y., Zhou G.J. & Leung K.M., 2017. Spatio-temporal variations in metal accumulation, RNA/DNA ratio and energy reserve in *Perna viridis* transplanted along a marine pollution gradient in Hong Kong. *Marine Pollution Bulletin*, 124: 736–742.
- Yusseppone M.S., Bianchib V.A., Castrob J.M., Noya Abada T., Minaberryc Y.S., Sabatinia S.E., Luquet C.M., Rios de Molinaa M.C. & Rocchetta I., 2020. In situ experiment to evaluate biochemical responses in the freshwater mussel *Diplodon chilensis* under anthropogenic eutrophication conditions. *Ecotoxicology and Environmental Safety*, 193: 110341.