

# Floristic and phytogeographic study of the vegetation of Djebel Médjounes (Setifian High Plains, Algeria)

Bariza Gourari<sup>1\*</sup>, Nacira Boulaacheb<sup>2</sup>, Vincent Andreu-Boussut<sup>2</sup> & Badr Eddine Belkhdja<sup>1</sup>

<sup>1</sup>Research Laboratory Urban Project, City Territory, Faculty of Nature and Life Sciences, University of Setif, Algeria

<sup>1</sup>Research Laboratory Urban Project, City Territory, Faculty of Medicine, University of Setif, Algeria

<sup>2</sup>Laboratoire ESO (UMR 6590 CNRS), Faculty of Geography, University of Le Mans, France

\*Corresponding author, e-mail: bariza.gourari.Etu@univ-lemans.fr

## ABSTRACT

The floristic and phytogeographical study of the pre-forest vegetation of Djebel Médjounes allowed for the evaluation of the floristic dynamics under anthropogenic and natural pressures due to climate change. Of the 237 phytosociological surveys, 420 taxa belonging to 53 families and 226 genera were recorded. The floristic composition reveals 186 Therophytes (45.03%), 133 Hemicryptophytes (32.20%), 33 Geophytes (8%), 40 Chameophytes (9.68%), 19 Nanophanerophytes (4.84%), and 2 Microphanerophytes, or 0.24% of the floristic cohort. The best represented families in terms of genera and species are Asteraceae (73 species, 17.33%) and Poaceae (55 species, 13.06%). The rest of the families have less than 50 species: Fabaceae (26 species, 11.16%), Brassicaceae (25 species, 5.93%), Lamiaceae (24 species, 5.7%), Caryophyllaceae (18 species, 4.27%), Boraginaceae (17 species, 4.03%), Apiaceae (13 species, 3.08%), Cistaceae and Crassulaceae (12 species, 2.85%), Ranunculaceae (10 species, 2.37%). From a phytogeographical perspective, the Mediterranean element is relatively predominant with 244 species (85.37%), of which 37 endemic species were inventoried. The identified floristic diversity component is induced by strong anthropogenic pressure (overgrazing, fires, exploitation) in addition to a regression of precipitation.

## KEY WORDS

Phytogeographical diversity; phytodiversity; anthropization; Djebel Medjounes; Algeria.

Received 21.11.2022; accepted 14.01.2023; published online 16.02.2023

## INTRODUCTION

Flora and vegetation are key elements for the study of biodiversity, their analysis allows for the evaluation of the state of ecosystems and helps in the development and implementation of conservation and protection strategies. The Mediterranean zone's forest heritage has been subjected to sustained decline for decades. Climate actions (summer

drought, irregular rain), anthropogenic (deforestation, overgrazing) and fires are the main causes of the extension of shrublands and the emergence of pre-forest vegetation groups. Such evolution has caused the substitution of mesophytic vegetation with xerophytic vegetation to varying degrees (Quezel & Barbero, 1990). Some environments such as rocky environments (as cliffs and rocks) play a major role in the conservation of certain plants populations (Quezel, 1957; Vogel et al., 1999).

Although much work has been done on Algerian flora and vegetation (Dahmani, 1984, 1997; Madoui, 1987, 1995, 2017, 2019; Merikhi, 1987, 1995; Gharzouli et al., 1989, 2000, 2002, 2005a, 2005b, 2007; Djebaili et al., 1989; Laouar, 1995; Khaloufi, 1995; Chermat, 1998; Quezel, 1999a, b; Boulaacheb et al., 2000, 2004, 2005, 2006a, 2006b, 2009, 2007, 2011; Bounar et al., 2001, 2003, 2008, 2011, 2012; Sarri et al., 2000, 2002; Vela et al., 2007; Rebbas et al., 2012; Yahy et al., 2012, Chermat et al., 2013, Gherzouli, 2013; Rebbas, 2014, etc.), the vegetation of Djebel Medjounes (High Plateaus of Sétif), located in the north, remains little known. Like the rest of the country, this region is subjected to a Mediterranean climate which influences the vegetation formations in general and the shrub formations in particular. Rainfall is centered on the cold seasons, and a long or very long summer drought, lasting 2 to 7 or 8 months per year Quezel (2000).

The High Plateaus of Sétif (Algeria) due to their great climatic, geological and geographical diversity are a refuge for many endemic and northern species. The floristic studies of these localities are an important source of information and their knowledge allows for the development of a program for

local, regional and national sustainable development and valorization that can be integrated into plans for the management and management of natural resources.

## MATERIAL AND METHODS

### Study area

The Djebel Medjounes (High Plateaus of Sétif) reaches an altitude of 1461 m and covers an area of 6,000 ha. It is located between the geographical coordinates  $36^{\circ}18'15''$  N Latitude and  $5^{\circ}31'17''$  E Longitude in DMS (degrees, minutes, seconds) or  $36.3042^{\circ}$  and  $5.5214^{\circ}$  (in decimal degrees) (Fig. 1).

The Djebel Medjounes is surrounded by a set of massifs, to the northeast the Djebels Moul el Djediane (1225 m) and Tamtarte (1008 m), to the northwest Djebel Chenatour (1370 m) and Djebel Senatour (1076 m); from East to West it is the Djebels Zerib (1317 m) and Zkarma (1361 m) and to the south the Djebels Mnaguer (1416 m) and Oulad Gues (1121 m).

The forest area is based on a mosaic of sedimentary formations from the Triassic, Cretaceous, Ter-




Figure 1. The study area's position in geographic terms (carried out by Gourari, 2020).

tiary, and Quaternary and are made up of an alternation of marls and calcareous marls (Vila, 1977, 1980). It is crisscrossed by numerous small alluvial and colluvial valleys. The hydrographic network consists of oueds such as oued Zatine, oued Dahab, oued Dahamcha, traversed by violent and frequent floods during the rainy season, are dry in summer.

The soils are calcimagnesian (rendzines, calcareous brown soils), raw and with little evolved minerals from erosion and input (M'Zoughem & Chenafa, 2006; Tedjar, 2011).

The study area receives 410 mm per year, the average calculated over a period of 32 years (1982–2020). The average monthly temperature is 15.48 °C, the coldest month is January with 2.03 °C, July is the hottest month with 33.67 °C. The dry season extends from May to September. According to the continentality index of Rivas-Martinez (1982, 2002), the climate is of the continental type (M-m = 31.64 °C), temperate-cool. Emberger's quotient (Q2) is equal to 51.48, with a semi-arid bioclimatic level with cold winters characterized by the concomitance of forest and steppe species (Le Houerou, 1995).

The study area is part of the mountain regions of the Tellian Atlas, the North African Mediter-

ranean domain (Quezel, 1978) also called "Mauritanian Mediterranean" according to LAPIE (1909–1914) and Maire (1926) or "Maghrebien Mediterranean" according to Barry et al. (1976). According to Meddour (2010) the study area belongs to the Maghrebo-Tellian domain, the Tello-Constantinian sector, and the Bibano Guelmois district.

### Data collection and floristic inventory

The floristic inventory was carried out based on 237 phytosociological surveys according to the sigmatist method of Braun-Blanquet (1932) and according to a subjective sampling taking into account the vegetation structure (matorrals, garrigues, grasslands, steppes, cliffs) and ecological factors (altitude, exposure, and slope).

The surveys were carried out in 14 stations, 7 on the north slope (Laamaera, Ouled Foudiya, Chorfa, Meliza, Siyab, Lahdada, and Boutouil) and 7 on the south slope (Lamaaz, Ballota, Rbiba, Ouled Ali Benacer, El Amria, Dajaria, and Sidi Halfaya) (Fig. 2).

During the years 2016–2018, plant samples were collected from 14 stations located on the north and south slopes of the Djebel Medjounes mountain. The surface area of the surveys ranged from




Figure 2. Location of the phytosociological surveys in Djebel Medjounes (Gourari & François, 2020).



10 m<sup>2</sup> for cliffs, 10–100 m<sup>2</sup> for rocky and barren grasslands, steppes, and 50–100 m<sup>2</sup> for shrublands. The plant samples were collected during all four seasons and the coordinates of the stations were recorded using GPS. The species samples were identified using the floras of Quezel & Santa (1962–1963) and Ozenda (1977) and the nomenclature used was based following Dobignard & Chatelain (2010–2013), Maire (1952–1987), and Chatelain et al. (2018). The classification of phytogeographic spectra was based on the work of Quezel & Santa (1962–1963), Le Floch et al. (1989), Le Houerou (1995), and Maire (1952–1987) and the biological types were defined according to the classification of Raunkiaer (1934). The rarity of species was based on the reference flora for Algeria (Quezel & Santa, 1962–1963).

## RESULTS AND DISCUSSIONS

### *Floristic analysis*

The inventory carried out using 237 surveys that listed 420 species belonging to 226 genera and 53 botanical families. The best represented families in terms of genera and species are Asteraceae (73 species and 47 genera, 17.33%) and Poaceae (55 species and 35 genera = 13.06%), followed by Fabaceae (26 species and 17 genera = 11.16%); Brassicaceae (25 species and 16 genera = 5.93%); Lamiaceae (24 species and 12 genera = 5.7%); Caryophyllaceae (18 species and 7 genera = 4.27%); Boraginaceae (17 species and 7 genera = 4.03%), Apiaceae (13 species and 8 genera = 3.08%); Cistaceae and Crassulaceae (12 species and 5 genera = 2.85%). The rest of the families have fewer than 10 species (Fig. 3). Asteraceae and Poaceae play a leading role on a global scale (Craven, 2009) and predominate in the Algerian flora (Quezel & Santa, 1962–1963). These results, whether for genera or species, are similar to those obtained by Gharzouli & Djellouli (2005), Bouнар (2001), Boulaacheb et al. (2005), Gharzouli (2007) and Chermat et al. (2013).

The report states that the ratio of number of families to number of species is 13%. The generic coefficient, or the ratio of number of genera to number of species, is 54%. Analysis of the biological spectrum (Fig. 4) shows the dominance of therophytes

(186 taxa = 45.03%). This richness in therophytes (*Guenthera amplexicaulis*, *Echinaria capitata*, *Calendula arvensis*, *Micropus supinus*, *Erodium cicutarium*, *Filago germanica*, *Bombycilaena erecta*, etc.) is a characteristic of arid Mediterranean areas where there is a high level of water stress (Daget, 1980; Madon & Medail, 1996). Hemicryptophytes represent 133 taxa, or 32.20%. Dahmani (1997), Barbero et al. (1988–2001), and Gharzouli (2007) report that their abundance in Maghreb countries is due to the richness of the soil in organic matter, altitude, and humidity (*Eryngium campestre*, *Thapsia garganica*, *Bupleurum spinosum*, *Eryngium triquetrum*, *Carduncellus pinnatus*, *Plantago lagopus*, *Teucrium polium* subsp. *capitatum*, etc.). Chamophytes are also well represented (40 species = 9.68%). According to Orshan et al. (1984) and Floret et al. (1990), chamophytes are fairly resistant to ecological and anthropogenic constraints and are able to adapt to drought (*Artemisia herba-alba*, *Launaea acanthoclada*, *Carlina gummifera*, *Helianthemum cinereum* subsp. *rotundifolium*, *Fumana thymifolia*, *Cistus creticus*, *Sedum album*, *Thymus munbyanus* subsp. *munbyanus*, *Teucrium pseudo-chamaepitys*, etc.).

The number of chamophytes may indicate a steppe influence, which could lead to a regression in regional preforest formations. Geophytes (33 taxa = 8%) are represented by Asphodelaceae (*Asphodelus ramosus*, *Asphodelus tenuifolius*), Poaceae (*Ampelodesmos mauritanicus*, *Echinaria capitata*, *Hordeum bulbosum*), and Orchidaceae (*Ophrys speculum*, *Ophrys lutea*). Nanophanerophytes are more or less represented, constituting 20 taxa (= 4.84%): *Crataegus monogyna*, *C. azarolus*, *Prunus dulcis*, *Quercus ilex* subsp. *ballota*, *Asparagus acutifolius*, *Pistacia terebinthus*, *P. lentiscus*, *Ziziphus lotus*, *Clematis flammula*, *Thymelaea hirsuta*, *Calicotome spinosa*, *Retama raetam*, *Genista tricuspidate*, *Daphne gnidium*, *Rhamnus lycioides* subsp. *oleoides*, *Globularia alypum*, *Rosmarinus eriocalyx*, *Prunus spinosa*).

Despite their low specific diversity, nanophanerophytes sometimes play a determining role in the establishment of a specific flora due to their coverage (Le Compte Barbet, 1975).

Microphanerophytes only have one taxon (*Cupressus sempervirens*), or 0.24% of the flora, this biological type, reforested in the region, seems to be better adapted to desertification.



The floristic composition is characterized by a high contribution (158 species = 38.82%) from the large distribution ensemble, which is a testament to the impact of human activity and the therophytisation of the flora (*Trifolium scabrum*, *Echinaria capitata*, *Geranium robertianum*, *Hippocrepis ciliata*, *Lathyrus sativus*, *Medicago minima*, *Vicia sativa*).

The cosmopolitan ensemble is well represented by 16 species (3.93%) and subcosmopolitan species count 2 species (0.49%). The species in this element have a very wide range of tolerance and are able to adapt to a wide range of ecological factors (*Avena sativa*, *Poa annua* subsp. *exilis*, *Sonchus arvensis*, *Sonchus oleraceus*, *Capsella bursa-pastoris*). They are mainly associated with moist environments, pastures, fields, and crops.

The transition ensemble consists of Mediterranean ensembles and neighboring ensembles. The largest group corresponds to Euro-Mediterranean species (48 species = 11.79%), including *Crataegus monogyna*, *Pallenis spinosa*, *Eryngium campestre*, *Fumana laevipes*, *Anisantha madritensis*, *Teucrium polium* subsp. *capitatum*, *Thapsia garganica*, etc. It is followed by the Eurasian element, which constitutes the major flora of temperate regions and plays an important role in the north of the Mediterranean. There are 30 species (7.37%) (*Torilis nodosa*, *Convolvulus arvensis*, *Geranium molle*, *Sedum album*, *Arenaria serpyllifolia*, etc.). The Mediterranean-Atlantic are 11 species (2.70%) (*Salvia lanigera*, *Ornithogalum umbellatum*, *Echinaria capitata*, *Centaurea aspera*, *Matthiola sinuata*, etc.).

The Mediterranean and Irano-Turanian group includes 8 species (1.96%), the rate of which is explained by a steppic influence, especially on the south slope (*Eremopyrum orientale*, *Aegilops geniculata* subsp. *geniculata*, *Hordeum bulbosum*, *Vulpia ciliata*, *Diptotaxis harra* subsp. *harra*, *Sisymbrium runcinatum*, *S. irio*). The presence of the Saharo-Sindian-Mediterranean (7 species = 1.71%) and Saharo-Mediterranean (3 species = 0.73%), *Micropus supinus*, *Erodium guttatum*, *Astragalus crenatus*, indicates the Saharan influence on this region. Their presence indicates the impact of climate change on the initial flora by favoring their installation.

The Euro-Siberian group is very poorly represented, with 4 species (0.98%): *Cichorium intybus*, *Carduus nutans*, *C. macrocephalus*, *Silene dioica*. Other categories are poorly represented with fewer than four species. The tropical ensemble is represented by 4 paleo-subtropical species (*Brachypodium distachyon*, *Gynandris sisyrrinchium*, *Anisantha rubens*, *Lolium rigidum*) and 2 subtropical species (*Muricaria prostrata* and *Matthiola lunata*). The endemic ensemble is present with 40 species, representing 9.50% of the total flora (Table 1): 28 North African endemics (6.65%), 3 Moroccan endemics (0.71%), 3 Algerian-Tunisian endemics (0.71%), 4 Algerian-Moroccan endemics (0.95%), and 2 Algerian endemics (0.47%).

There are 40 rare species, of which 21 (2.47%) are very rare (R), 10 (6.5%) are rare (AR), and 9 (3.1%) are fairly rare (RR) (Fig. 5). Of the 40 species, 8 are on the list of non-cultivated plant




Figure 4. Rates of biological types identified in Djebel Medjounes.




Figure 5. Attendance rate of species identified in Djebel Medjounes.

species protected by the Algerian law of January 4, 2012 (*Tragopogon porrifolius* subsp. *macrocephalus*, *Silene colorata* subsp. *amphorina*, *Lomelosia stellata*, *Thymus guyonii*, *Ononis natrix* subsp. *angustissima*, *Linum grandiflorum*, *Salix triandra*, *Linaria pelisseriana*).

Boulaacheb (2009) notes that ruderal, nitrophilic, and unpalatable species are indicators of disturbed and open habitats, signs of strong human activity. These include: *Ornithogalum umbellatum*, *Asphodelus ramosus*, *Asphodelus tenuifolius*, *Leontodon hispidus*, *Eryngium campestre*, *Eryngium bourgatii*, *Eryngium triquetrum*, *Sixalix atrop-*

*urpurea*, *Galium odoratum*, *Galium fruticosum*, *Salvia lanigera*, *Salviaverbenaca*, *Plantago ser-raria*, *Plantago lanceolata*, *Plantago coronopus*, *Hypochaeris achyrophorus*, *Helianthemum nummularium*, *Helianthemum cinereum*, *Rumex crispus*, *Rumex bucephalophorus* subsp. *gallicus* f., *Trifolium stellatum*, *Trifolium glomeratum*, *Centaurea oranensis*, *Centaurea sphaerocephala*, *Minuartia montana*, *Centaureum pulchellum*, *Centaureum erythraea*.

On the south flank of the Djebel Medjounes mountain, we also observed the infiltration of steppe and Saharan plants, indicating the impact of

Species	Chorology	Species	Chorology
<i>Astragalus froedinii</i> Murb.	End. du Maroc	<i>Silene patula</i> subsp. <i>amurensis</i> (Pomel) Jeanmonod	End. N. A.
<i>Bunium fontanesii</i> (Pers.) Maire	End. N. A.	<i>Carduncellus pinnatus</i> (Desf.) DC.	End. N. A.
<i>Sedum pubescens</i> Vahl	End. N. A.	<i>Thymus guyonii</i> de Noé	End. Alg.
<i>Silene atlantica</i> Coss. & Durieu	End. N. A.	<i>Cirsium palustre</i> (L.) Scop.	End. N. A.
<i>Helianthemum ruficomum</i> (Viv.) Spreng.	End. N. A.	<i>Atractylis caespitosa</i> Desf.	End. N. A.
<i>Herniaria hirsuta</i> L.	End. N. A.	<i>Leontodon hispidus</i> L.	End. N. A.
<i>Echium horridum</i> Batt.	End. N. A.	<i>Echium parviflorum</i> Moench	End. N.A.
<i>Echium humile</i> Desf.	End. N. A.	<i>Echium humile</i> subsp. <i>pyncnanthum</i> (Pomel) Greuter & Burdet	End. N.A.
<i>Paronychia arabica</i> (L.) DC.	End. N. A.	<i>Origanum vulgare</i> subsp. <i>glandulosum</i> (Desf.) Ietsw.	End. N. A.
<i>Galium tricornutum</i> Dandy	End. N.A.	<i>Iberis odorata</i> L.	End. Alg. Mar.
<i>Diploxys simplex</i> (Viv.) Spreng.	End. N. A.	<i>Rhaponticum acaule</i> (L.) DC.	End. N. A.
<i>Hertia cheirifolia</i> (L.) Kuntze	End. N. A.	<i>Delphinium pentagynum</i> Lam.	End. Alg. Tun
<i>Hedysarum boveanum</i> Bunge ex Basiner	End. N. A.	<i>Rosmarinus eriocalyx</i> Jord. & Fourr.	End. N. A.
<i>Astragalus armatus</i> Willd. subsp. <i>armatus</i>	End. N. A.	<i>Genista tricuspida</i> Desf.	End. N. A.
<i>Thymus munbyanus</i> Boiss. & Reut. subsp. <i>munbyanus</i>	End. N. A.	<i>Gagea granatellii</i> (Parl.) Parl.	End. Alg.
<i>Thymus broussonetii</i> Boiss. subsp. <i>broussonetii</i>	End. du Maroc	<i>Hedysarum pallidum</i> Desf.	End. N.A.
<i>Thymus munbyanus</i> subsp. <i>ciliatus</i> (Desf.) Greuter & Burdet	End. N. A.	<i>Linum usitatissimum</i> L.	End. Alg. Mar.
<i>Hippocrepis atlantica</i> Ball	End. N. A.	<i>Linum bienne</i> Mill.	End. Alg. Mar.
<i>Sinapis pubescens</i> L.	End. Alg. Tun	<i>Vulpia alopecuros</i> (Schousboe) Dumort.	End. du Maroc
<i>Tragopogon porrifolius</i> L. subsp. <i>macrocephalus</i> Batt.	End. Alg. Mar.	<i>Linaria reflexa</i> (L.) Chaz. subsp. <i>brevicalcarata</i> D.A.Sutton	End. Alg. Tun.

Table 1. List of endemic species (End.) of North Africa, or limited to some of its regions, found in Jebel Medjounes.



climate change on one hand and over exploitation on the other hand: *Artemisia herba alba*, *Ajuga iva*, *Thymelaea hirsuta*, *Ziziphus lotus*, *Retama raetam*, *Stipa parviflora*, *Macrochloa tenacissima*, *Lygeum spartum*, *Anisantha rubens*, *Atractylis serratoloides*, *Lobularia canariensis* subsp. *rosula-venti*, *Erodium guttatum*, *Helianthemum ruficomum*, *Reseda arabica*, etc.

On the south slope, with a very high population density, the vegetation is characterized by herbaceous grasslands and rocky areas that provide information about the type of activity (pastoralism) and the degree of openness of the environment (Fig. 6).

On the contrary, the north flank with very low population density is more or less preserved. It is marked by the presence of woody plants and is characterized by matorrals with *Rhamnus alaternus* and matorrals with holm oak (Figs. 7, 8). Both slopes are characterized by the presence of Diss (*Ampelodesmos mauritanicus*) and *Calicotome spinosa*. These species result from the degradation of wooded matorrals with green oaks (Boulaacheb, 2005).

Mining in the north slope (Boutouil, Lahdada) is known by ecologists as a form of nature aggression in all its forms (Fig. 9).



Figure 6. Clear shrublands. South slope of ouled Ali Benaser.



Figure 7. Matorrals with gaps. North slope of Mliza.



## CONCLUSIONS

The pre-forest vegetation of Djebel Médjounes is characterized by a very diverse flora with 420 species of which 40 endemic taxa. A significant number of species with medicinal value (63 species) could improve the income of local populations while ensuring the conservation of floral diversity. The biological spectrum generally corresponds to a characteristic pattern of Mediterranean pre-forest formations with the dominance of therophytes related to an increasing gradient of aridity in the northeastern Algerian High Plateaus.

The vegetation cover of this massif shows a strong contrast between the north slope where shrublands and garrigues dominate and degraded forms of shrublands and the south slope where stripped lawns formed of Saharan and steppe species dominate. Climate change and human pressure (overgrazing, harvesting) are the main causes of degradation of the Djebel Médjounes, causing a decline in species and a reduction in plant diversity.

This work on the plant diversity of the Djebel Médjounes provides a scientific basis for planning sustainable development linked to the preservation of this ecosystem and its heritage.



Figure 8. Matorrals. North slope of Laamara.



Figure 9. Mining operation in the north slope (Boutouil).

## REFERENCES

- African Plants Database, 2018. African Plants Database (version 3.4.0). Conservatoire et Jardin botaniques de la Ville de Genève and South African National Biodiversity Institute, Pretoria.  
<http://www.ville-ge.ch/musinfo/bd/cjb/africa/>
- Barbero M., Bonin G., Loisel R. & Quézel P., 1990. Changes and disturbance of forest ecosystems caused by human activities in the western part of the Mediterranean basin. *Vegetatio*, 87: 151–173.  
<https://doi.org/10.1007/BF00042952>
- Barbero M. & Quézel P., 1995. Desertification, aridification in the mediterranean region and global change. In: Bellan D., Bonin G. & Emig C. (Eds.), *Functioning and dynamics of natural and perturbed ecosystems*, Paris, Lavoisier, Intercept Ltd., pp. 549–569.
- Barbero M., Loisel R., Médail F. & Quézel P., 2001. Signification biogéographique et biodiversité des forêts du bassin méditerranéen. *Bocconea*, 13: 11–25.
- Barbero M., Quézel P. & Rivas-Martinez S., 1981. Contribution à l'étude des groupements forestiers et préforestiers du Maroc. *Fitosociologia*, 9: 311–412.
- Boulaaheb N., 2009. Etude de la végétation terrestre et aquatique du djebel Megriss (Nord Tellien, Algérie). Analyse floristique, phytosociologique et pastorale. Thèse Doctorat science Université F. Abbas, Sétif, 331 pp.
- Boulaaheb N., Gharzouli R. & Djellouli Y., 2005. Approche phytosociologique du Djebel Megriss (nord de Sétif, Algérie). *Bulletin de la Société Botanique du Centre-Ouest, Nouvelle série*, 6: 345–362.
- Bouhar R., 2001. Etude phytoécologique, cartographie et aménagement du Massif des Babors. Mémoire de Magister Université F. Abbas, Sétif, 110 pp.
- Bagnouls F. & Gaussen H., 1953. Saison sèche et indice xérothermique. Doc.: Cartes Product. veget. Serie: Generalites, 3 (193–239), art. 8, 47 pp. + 1 carte.
- Barry J.P., Celles J.C. & Faurel L., 1974. Carte internationale du tapis végétal et des conditions écologiques. Feuille d'Alger au 1/1.000.000e. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, Alger, 164 pp.
- Chermat S., Djellouli Y. & Gharzouli R., 2013. Dynamique régressive de la végétation des hautes plaines sétifiennes: érosion de la diversité floristique du Djebel Youssef (Algérie). *Revue d'écologie - La terre et la vie*, 68: 85–100
- Chatelain C., Medjahdi B. & Benhouhou S., 2018. La flore du Magreb 2018 basé sur la "Nouvelle flore d'Algérie et des régions désertiques méridionales" de Quézel et Santa 1963. version 2.2sept.2018  
<https://efloramaghreb.org/>
- Craven P., 2009. Phytogeographic study of the Kaokoveld centre of endemism. Dissertation presented for the degree of Doctor of Philosophy at Stellenbosch University, 234 pp.
- Daget Ph., 1980. Sur les types biologiques botaniques entant que stratégie adaptative (cas des thérophytes). In: *Recherches d'écologie théorique: les stratégies adaptatives*. Paris, pp. 89–114.
- Djellouli Y., 1990. Flore et climat en Algérie septentrionale. Déterminismes climatiques de la répartition des plantes. Thèse de Doctorat, Université des sciences et de la technologie Houari Boumédiène, Alger.
- Gharzouli R. & Djellouli Y., 2005. Diversité floristique de la Kabylie des Babors (Algérie). *Sécheresse*, 16: 217–223.
- Dobignard A. & Chatelain C., 2018. Index synonymique de la Flore Afrique du Nord, vol. 1–5. Edit. Conservatoire et Jardin botaniques de la Ville de Genève, Éditions. Des Conservatoire et Jardin Botaniques, Genève, 455 pp.
- Dobignard A. & Chatelain C., 2010. Index synonymique de la Flore d'Afrique du Nord, Vol. 1 (Pteridophyta, Gymnospermae, Monocotyledoneae). Ville de Genève, Éditions Des Conservatoire et Jardin Botaniques, Genève, 455 pp.
- Emberger L., 1955. Une classification biogéographique des climats. *Recueil des travaux des Laboratoires de Botanique, Géologie et Zoologie de la Faculté des Sciences de l'Université de Montpellier*, 7: 1–43.
- Emberger L., 1952. Sur le quotient pluviothermique. *Comptes rendus de l'Académie des Sciences Paris*, 23: 2308–2310.
- Ellenberg I. & Mueller-Doubois D., 1967. A key to Raunkier plant life forms with revised subdivisions. *Berichte des Geobotanischen Institutes der Eidgen. Technischen Hochschule, Stiftung Rübel*, 378: 56–63.
- Floret C., Galan M.J., Le Floch E. & Romane F., 1990. Growth forms and phenomorphology traits along an environmental gradient: tools for study vegetation? *Journal of Vegetation Science*, 1: 71–80.
- Gharzouli R. & Djellouli Y., 2005. Diversité floristique de la Kabylie des Babors (Algérie). *Sécheresse*, 16: 217–223.
- Gharzouli R., 1989. Contribution à l'étude de la végétation de la chaîne des Babors. Analyse phytosociologique des Djebels Babor et Tababort. Mémoire de Magister Institute Nationale Ens. Supérieure de Sétif, 235 pp.
- Gharzouli R., 2007. Flore et végétation de la Kabylie des Babors. Etude floristique et phytosociologique des groupements forestiers et post-forestiers des djebels Takoucht, Adrar ou Melal, Tababort et Babor. Thèse Doctorat en Biologie Végétale Université Ferhat Abbas de Sétif, 387 pp.
- Gharzouli C., 2013. Anthropisation et dynamique des

- zones humides dans le nord-est algérien: Apport des études palynologiques pour une gestion conservatoire. Thèse Doctorat Science Université de Toulouse, 208 pp.
- Khaloufi-Souici N., 1995. Contribution à l'étude de la végétation du Tell Sétifien. Analyse phytosociologique des Djebels Tafat, Anini et Megriss. Mémoire Magister Université F. Abbas Sétif, 148 pp.
- Le Compte-Barbet O., 1975. Introduction à une étude de l'endémisme végétal au Maroc. Travaux de recherche au Centre national de la recherche scientifique, 249: 15-46.
- Le Floch E., Akka O., Hirmiz A.K.T., Masri A., Meziani K. & Tadros K., 1989. Les techniques de développement pastoral: plantation d'arbustes fourragers. Vol 3. FAO, 204 pp.
- Le Houerou H.N., Claudin J. & Pouget M., 1977. Étude bioclimatique des steppes algériennes. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord, n.s., 68: 33-70.
- Le Houerou H-N., 1980. L'impact de l'homme et de ses animaux sur la forêt méditerranéenne. Forêt Méditerranéenne, 2: 155-174.
- Le Houerou H-N., 1995. Bioclimatologie et biogéographie des steppes arides du Nord de l'Afrique: diversité biologique, développement durable et désertisation. In: Le Houerou H.-N. (Ed.), Bioclimatologie et biogéographie des steppes arides du Nord de l'Afrique : diversité biologique, développement durable et désertisation. Montpellier: CIHEAM, 1995, pp. 1-396. (Options Méditerranéennes: Série B. Etudes et Recherches; n. 10.
- Lapie G., 1914. Aperçu phytogéographique sur la Kabylie des Babors. Revue générale de botanique, Vol. G Bonnier, pp. 417-424.
- Meddour R., 2010. Bioclimatologie, phytogéographie et phytosociologie en Algérie. Thèse de Doctorat, université Mouloud Mammeri, TiziOuzou, Algérie, 461 pp.
- M'Zoughem K. & Chenafa W., 2006. Etude géotechnique de la stabilité des talus dans la carrière d'Ain El Kebira (Sétif). Mémoire de fin d'étude en vue de l'obtention du diplôme d'Ingénieur d'Etat en Géologie. UFAS Sétif, 68 pp.
- Merikhi R., 1987. Contribution à la connaissance de la cédraie du massif du Boutaleb. Mémoire de Université F. Abbas, Sétif, 21 pp.
- Merikhi R., 1995. Contribution à l'étude la végétation de monts du Hodna; étude phytosociologique du massif du Boutaleb. Thèse de Magister, Université F. Abbas, Sétif, 179 pp.
- Maire R., 1926. Carte phytogéographique de l'Algérie et de la Tunisie. 1/1 500000, Gouvernement général d'Alger, Service cartographie, Alger.
- Maire R., 1928. Origine de la flore des montagnes de l'Afrique du Nord. Mémoires - Société de Biogéographie, 2: 187-194.
- Maire R., 1952-1987. Flore de l'Afrique du Nord (Maroc, Algérie, Tunisie, Tripolitaine, Cyrénaïque, Sahara). Vol. i-xvi. Lechevalier, Paris.
- Madon O. & Médail F., 1997. The ecological significance of annuals on a Mediterranean grassland (Mont Ventoux, France). Plant Ecology, 129: 189-199.
- Ozenda, P. 1977. Flore du Sahara. 2ème éd., CNRS, Paris, 622 pp.
- Orshan G., Montenegro G., Avila G., Aljaro M.E., Walckowiaak A. & Mujica A.M., 1984. Plant growth forms of Chilean matoral. A monocharacter growth form analysis along an altitudinal transect from sea level to 2000 m a.s.l. Bulletin de la Société botanique de France, 131: 411-425.
- Quézel P., 1978. Analysis of the flora of Mediterranean and Saharan Africa. Annals of the Missouri Botanical Garden, 65: 479-537.
- Quezel P., 1964. L'endémisme dans la flore de l'Algérie. Comptes rendus de la Société de Biogéographie, 361: 137-149.
- Quezel P., 1999a. Réflexions sur l'évolution de la flore et de la végétation au Maghreb méditerranéen. Ibis Press, Paris, 117 pp.
- Quezel P., 1999b. Les grandes structures de végétation en région méditerranéenne : facteurs déterminants dans leur mise en place post-glaciaire. Geobios 32: 19-32.  
[https://doi.org/10.1016/S0016-6995\(99\)80081-3](https://doi.org/10.1016/S0016-6995(99)80081-3).
- Quezel P., 2000. Réflexion sur l'évolution de la flore et de la végétation au Maghreb méditerranéen. Ibis Press Paris, 117 pp.
- Quezel P. & Barbero M., 1990. Les forêts méditerranéennes problèmes posés par leur signification historique, écologique et leur conservation. Acta Botanica Malacianata, 15: 145-178.  
<https://doi.org/10.24310/abm.v15i.9300>
- Quezel P. & Médail F., 2003. Écologie et biogéographie des forêts du bassin méditerranéen. Elsevier, Paris, 573 pp.
- Quezel P. & Santa S., 1962-1963. Nouvelle flore de l'Algérie et des régions désertiques méridionales. CNRS, Paris. 2 vol., 1170 pp.
- Raunkiaer C., 1934. The life form of plants and statistical plant geography. Collected papers, Clarendon Press Oxford, 623 pp.
- Rebbas K., 2014. Développement durable au sein des aires protégées algériennes, cas du Parc National de Gouraya et des sites d'intérêt biologique et écologique de la région de Béjaïa. Thèse Doctorat en Biologie Végétale Université Ferhat Abbas de Sétif, 192 pp.
- Rebbas K., Bounar R., Gharzouli R., Ramdani M., Djelouli Y. & Alatu D., 2012. Plantes d'intérêt médi-



- nale et écologique dans la région d'Ouanougha (M'Sila, Algérie). *Phytothérapie*, 10: 131–142.  
<https://doi.org/10.1007/s10298-012-0701-6>
- Rivas-Martínez S., Díaz T.E., Fernández-González F., Izco J., Loidi X., Lousã M. & Penas A., 2002. Vascular plant communities of Spain and Portugal: addenda to the syntaxonomical checklist of 2001. *Itinera Geobotanica*, 15: 5–922.
- Rivas-Martínez S., 1982. Etages bioclimatiques, secteurs chronologiques et séries de végétation de l'Espagne méditerranéenne. *Ecologia Mediterranea*, 8: 275–299.
- Rivas-Martínez S. & Penas Merino Á. (Coord. Scient), 2003. Atlas manual de los habitats de España. Ministerio de Medio ambiente. Secretaria general de Medio ambiente. Direccion General de Conservacion de la Natureza, Tragsa, 487 pp.
- Rivas-Martínez S., 1982. Etages bioclimatiques, secteurs chronologiques et séries de végétation de l'Espagne méditerranéenne. *Ecologia Mediterranea* 8: 275–287.
- Sadki N., 1988. Contribution à l'étude des groupements à olivier et lentisque de la région d'Annaba. Essai phytosociologique. Thèse Doctorat 3<sup>ème</sup> cycle. U.S.T.H.B., Alger, 213 pp.
- Tedjar L., 2012. Impact des rejets (unités industrielles: cimenterie et BCR) sur l'environnement dans la région d'Ain EL Kebira (Sétif). Thèse Doctorat en Biologie Animal Université Ferhat Abbas, Sétif, pp. 13–50.
- Vila J.M., 1974. Le Rocher de Constantine, stratigraphie, microfaunes et position structurale. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 65: 385–392.
- Vila J.M., 1977. Carte géologique de l'Algérie au 1/50 000, feuille N° 74, El Aria avec notice explicative détaillée (levés de S. Guellal). Service de la carte géologique de l'Algérie, SONATMCH.
- Vila J.M., 1980. La chaîne alpine d'Algérie orientale et des confins algéro-tunisiens. Thèse Sciences Université Pierre et Marie Curie, Paris 6, 3 vol., 663 pp.
- Zohary D. & Hoff M., 1974. Domestication of pulses in the old world. *Sciences*, 182: 887–894.