

Rare and endangered species of Shahdag National Park (Azerbaijan) with special status

Sayyara Jamsid Ibadullayeva¹, Elman Farhad Yusifov¹ & Arzu Bahadur Mustafayev²

¹Institute of Botany, Ministry of Science and Education of the Republic of Azerbaijan

²State Environmental Security Service, Ministry of Ecology and Natural Resources of Azerbaijan Republic; e-mail: yusifov_e@yahoo.com; atani@bk.ru

Corresponding author, e-mail: sayyara.ibadullayeva@mail.ru

ABSTRACT

This work presents information about the geographical ranges of the vegetation of the Shahdag National Park (NP) belonging to the Greater Caucasus (within the borders of Azerbaijan) range, the current status of endemic, relict and rare species. The geographic-areological analysis of the region shows that the flora of the Shahdag area is formed by the predominance of Boreal and Xerophile types. From the studies, it was found that the spread of endemic taxa in NP to neighboring areas reduced them to the subendemic level. In total, 7 species of Azerbaijan endemic and 88 subendemic species are distributed in the territory of Shahdag NP, of which 69 are described only from the Caucasus and are considered endemic to the Caucasus, while others are distributed as far as the borders of Dagestan, Turkey and Iran. In terms of determining the development history of the local flora, 24 relict species have been determined. In general, 14 species were evaluated as vulnerable (VU), 10 species as Near Threatened (NT), and 5 species as endangered (EN) in the Shahdag area.

KEY WORDS

National Park; endemic; relict; rare species; geographical range.

Received 08.05.2023; accepted 04.08.2023; published online 05.09.2023

INTRODUCTION

Without studying the flora of certain territories, it is difficult to ensure the protection of its plants that have a special status, their rational use, and the implementation of improvement measures in territories affected by the negative impact of environmental, anthropogenic and zoogenic factors. Also in botanical geography, the study of the endemic specie is of great importance. The endemic or relict species distributed in a certain area is the main indicator of the specificity and originality of that flora. As a result of studying the composition of endemic taxa, the regularity of flora formation,

phylogenetic condition, chorological and eco-phytocenological characteristics, as well as genesis, become clear. Endemics are understood as taxa the distribution of which is limited to a certain territory (Tolmachev, 1986). The geographic distribution of endemic taxa may vary. Species, the range of which goes beyond the boundaries of the area under study and is not limited to a certain area, are called subendemic. Endemic and relict plants of the vegetation of Azerbaijan and the Caucasus have been studied by many authors (Askerov, 2006–2008; Salimov et al., 2021; Babakishiyeva, Ibadullayeva, 2021). Information on 240 endemic species for the flora of Azerbaijan has been provided by Akhundov

(1973). Subsequently, Grossheim (1939) recorded the presence of 1153 endemic species in Caucasus and Asgarov (2016) provided information on 146 endemic and 402 subendemic plants that make up the gene pool of the flora of Azerbaijan. As a result of the research towards the study of the endemic plants of Azerbaijan in recent years, some of the endemic species have been assessed as subendemic (Yusifov et al., 2022).

Floristic research is extremely important in terms of studying the vegetation of the region. As a result of the analysis of the research conducted in this direction and the identification of the plant samples obtained during the expeditions, the flora of the territory of the Shahdag National Park has been fully studied and systematically analyzed (Ibadullayeva et al., 2014). During the research period, a schematic description of higher plants consisting of 1603 species in the flora of the Shahdag National Park area has been developed. It was determined that 31 species of spore and 1572 species of seed (12 species of gymnosperms, 1560 species of angiosperms, including 294 species of monocotyledons, 1266 species of dicotyledons) are distributed in the local flora (Mustafayev, 2019).

In recent years, in order to protect the biodiversity of rare species that have a limited range and are on the verge of extinction, extensive research work is being carried out (Musayev, 2005; Talibov et al., 2010; Mursel et al., 2019; Ibadullayeva et al., 2021; Mehdiyeva et al., 2022). The assessment of rare and endangered species is first carried out at the regional level, then at the national level, and finally at the international level. For this purpose, the International Union for Conservation of Nature (IUCN) was developed, and in Azerbaijan, according to the criteria of the Red List of this Union, the protection status of rare species was determined (IUCN, 2001).

Taking all this into account, the goal was to study the origin of plants with a special status of the Shahdag NP, including the geographic ranges and the reasons for their decline.

MATERIAL AND METHODS

During the research, attention was paid to the

latest nomenclatures data following Portenier (2000) and its methods based on phytochorions in botanical-geographical zoning. The flora of the region was geographically analyzed using the works of Grossheim (1939; 1939–1967). The protection status of rare species is given according to the IUCN “Red Data Book” (2003) criteria and the “Red Books” of the Republic of Azerbaijan (1989, 2013, 2023). In addition, attention was paid to the publications on rare and endemic species (Asgarov, 2016; Ibadullayeva et al., 2021).

RESULTS AND DISCUSSION

Floristic geographical studies, which provide information on the geographic location of the intraspecific smallest taxa and the center of distribution of rare species (relict, endemic), play a key role in the protection of vegetation, the genetic origin of species, on the physical-geographical conditions, modern and past, of the area where it is inhabited, for determining the geographical element and the areal types to which they belong, and determining the distribution and centers of formation of the species.

Based on the herbarium materials and literature sources collected during the research years, for the flora of Shahdag area, zonal areal type were determined. The types, classes and groups of all species distributed in the area's flora according to their geographical range are defined and reflected in Table 1. The geographic analysis of the region shows that the flora of the Shahdag is formed by the predominance of boreal and xerophilous types. Species with xerophilic areal type constitute almost the majority of the region flora with 571 species (35.5%), the boreal areal type with 538 species (33.6%) and the Caucasian areal type with 329 species (20.5%). Steppe areal type is represented by 48 species (3%). Indeterminate areal type is represented by 41 species (2.5%), ancient areal type by 24 species (1.5%), adventive areal type by 21 species (1.3%), desert areal type by 17 species (1.1%), and cosmopolitan areal type by 14 species (0.9%).

There are 24 species - *Anizantha tectorum* (L.) Nevski, *Chenopodium foliosum* Aschers, *Ch. vulvaria* L., *Krascheninnikovia ceratoides* (L.)

№	Areal types	Species	%	Classes	%	Groups	%
1	Ancient	24	1.5	2	13.3	6	14
2	Boreal	538	33.6	3	20	7	16.3
3	Steppe	48	3	3	20	8	18.6
4	Xerophile	571	35.6	3	20	16	37.2
5	Desert	17	1.1	2	13.3	3	7
6	Caucasus	329	20.5	1	6.7	2	4.7
7	Adventive	21	1.3	1	6.7	1	2.2
8	Cosmopolitan	14	0.9	-	-	-	-
9	Undeterminate	41	2.5	-	-	-	-
Total		1603	100	15	100	43	100

Table 1. Composition of species in the flora of Shahdag NP according to geographical areal types, classes and groups.

Guelldents., *Rumex scutatus* L., *Platanus orientalis* L., *Juglans regia* L., *Viola sicheana* W.Beck., *Datisca cannabina* L., *Thymelaea passerina* (L.) Coss. et Germ., etc - of relict plants in the ancient areal type in the area.

There are 9 main and 7 transition groups in 3 classes of the xerophilic areal type (Mediterranean, Western Asia, Central Asia) represented by more species. Most species are distributed in Western Asia (69 species), Mediterranean (74 species), Irano-Turanian (53 species), Azerbaijan endemics (38 species), Iran (34 species) and Mediterranean-Irano-Turanian (33 species) groups. Turanian group of desert areal type is represented by the species *Lonicera iberica* Bieb., *Viburnum opulus* L., *Adonis bienertii* Butk., Eastern Transcaucasian group by the species *Nonnea rosea* (Bieb.) Link, *Acer laetum* C.A.Mey., *A. hyrcanum* Fisch. et C.A.Mey., *Lonicera iberica* Bieb., *Papaver arenarium* Bieb., *Melandrium latifolium* (Poir.) Maire, *Sclerochloa dura* (L.) Beauv, Sacharo-Iranian group by the species *Lepidium sativum* L. The Caucasian areal type includes 329 species in one class and 2 groups, of which *Callicephalus nitens*, *Xeranthemum longepapposum*, *X. cylindraceum*, *Centaurea behen*, *Achillea tenuifolia* species are more common. Cosmopolitan areal type is represented by only 41 species - *Cerastium glomeratum* Thuill., *Triglochin*

palustre L., *Botrychium lunaria* (L.) Sw., *Ophioglossum vulgatum* L., *Poa palustris* L., *P. caucasica* Trin., *P. annua* L., *Sonchus oleraceus* L., *Veronica biloba* Schreb., *Lemna minor* L., *L. trisulea* L., etc.

The species *Sisymbrium altissimum* belonging to steppe areal type of the region is included in Pontic-Sarmatian group, *Stipa lessingiana* species to Pannon-Sarmatian, *Phlomis pungens* species to Eastern Mediterranean-Pontic transition group, *Eremopyrum triticeum*, *Catabrosella humilis*, *Ceratocarpus arenarius*, *Atriplex cana* species to Sarmatian main group.

In the boreal areal type, which takes the second place by the number of species, the main place is occupied by Palearctic with 142 species and European areal elements with 176 species. In the flora of the region, the Holarctic is represented by 87 species, and the Western Palearctic by 65 species.

Thus, the vegetation of the Shahdag area was formed due to the Western Asia, the Mediterranean Sea, Irano, Turanian xerophilic elements, on the one hand, and Palearctic, European, Holarctic boreal elements on the other hand. Local oborogenic species with Atropathan and Caucasian elements also play an important role in the formation of the area's flora.

Endemic and relict plants belonging to various floristic elements are found in the composition of

Subendemics	<i>Galium brachyphyllum</i> Roem. et Schult.	Caucasus, Iran, Turkey
	<i>Quercus macranthera</i> Fisch. et C.A. Mey. ex Hohen.	Caucasus
	<i>Malabaila sulcata</i> Boiss.	Caucasus
	<i>Merendera eichleri</i> (Regel) Boiss.	Caucasus
	<i>Medicago caucasica</i> Vass.	Caucasus
	<i>M. glutinosa</i> Bieb.	Caucasus
	<i>Nonea alpestris</i> (Stev.) G. Don. f.	Caucasus, Iran
	<i>Nepeta supina</i> Stev.	Caucasus
	<i>N. cyanea</i> Stev.	Caucasus, Turkey
	<i>Ornithogalum sintenisii</i> Freyn	Caucasus
	<i>Onosma levinii</i> T.N. Pop.	Caucasus, Turkey
	<i>Pimpinella aromatica</i> Bieb.	Caucasus, Iran
	<i>Poa meyeri</i> Trin. ex Roshev.	Caucasus
	<i>Pseudovesicaria digitata</i> (C.A.Mey.) Rupr.	Caucasus, Iran, Turkey
	<i>Pulsatilla albana</i> (Stev.) Bercht. et J. Presl.	Caucasus, Turkey
	<i>Ranunculus crassifolius</i> (Rupr.) Grossh.	Caucasus
	<i>Rosa alexeenkoi</i> Crep. ex Juz.	Caucasus
	<i>R. sachokiana</i> P. Jarosch.	Caucasus
	<i>R. sosnowskyi</i> Chrshan.	Caucasus
	<i>R. komarovii</i> Sosn.	Caucasus
	<i>Salix kuznetzowii</i> Laksch. ex Goerz	Caucasus
	<i>Salvia verbascifolia</i> Bieb.	Caucasus
	<i>Saxifraga pseudolaevis</i> Oetting.	Caucasus
	<i>Scutellaria oreophila</i> Grossh.	Caucasus
	<i>Silene caucasica</i> Boiss.	Caucasus
	<i>S. lacera</i> (Stev.) Sims	Caucasus
	<i>Symphyloloma graveolens</i> C.A. Mey.	Caucasus
	<i>Senecio grandidentatus</i> Ledeb.	Caucasus
	<i>S. kubensis</i> Grossh.	Caucasus
	<i>Serratula caucasica</i> Boiss.	Caucasus
	<i>Sedum obtusifolium</i> C.A. Mey.	Caucasus
	<i>S. subulatum</i> (C.A. Mey.) Boiss.	Caucasus
	<i>S. stevenianum</i> Rouy et Camus	Caucasus, Iran
	<i>Stemmacantha pulchra</i> (Fisch. et C.A. Mey.) Dittrich	Caucasus, Turkey
	<i>Sorbus caucasica</i> Zinserl.	Caucasus
	<i>Scrophularia variegata</i> Bieb.	Caucasus
	<i>Stachys pauli</i> Grossh.	Caucasus
	<i>Trigonocaryum involucratum</i> (Stev.) Kusn.	Caucasus
	<i>Tanacetum meyerianum</i> Sosn. (<i>Pyrethrum meyerianum</i> Sosn.)	Caucasus
	<i>Valerianella amblyotis</i> Fisch. et C.A. Mey.	Caucasus, Iran
	<i>Valeriana cardamines</i> Bieb.	Caucasus
	<i>Veronica peduncularis</i> Bieb.	Caucasus, Iran
	<i>V. petraea</i> (Bieb.) Stev.	Caucasus
	<i>V. minuta</i> C.A.Mey.	Caucasus
	<i>Vicia loiseleurii</i> (Bieb.) Litv.	Caucasus, Iran
	<i>Vavilovia formosa</i> (Stev.) Fed.	Caucasus, Iran, Turkey

Table 2. Endemic and subendemic species of flora of Shahdag NP.

vegetation of Shahdag area. Previously, there was information about the existence of 277 species of endemic plants in the flora of the territory (Flora of Azerbaijan, 1950-1961), of which 252 species (91%) are endemic to the Caucasus, and 25 species

endemic to Azerbaijan. This makes up 17.28% of the area's flora.

Endemism can be characterized by a relatively small area. Among the endemics of Azerbaijan, elements of the Caucasian group dominate with 9

species, elements of Palearctic, Iranian-Turanian, Atropatan and Mediterranean with 2 species each, elements of adventive, Armenian, Holarctic, Asia Minor, Pontic, Caucasus-Central Asia, North Iranian are very few (1 species each), and the range of one species is unknown. They are mainly distributed in the forest, forest edge cenoses, subalpine, alpine and subnival zones of Shahdag NP. Caucasian endemics are distributed on the Caucasus, Western Asia, Asia Minor, Palearctic, Atropathan, Irano, Mediterranean, and European elements.

The analysis of recent studies towards the analysis of the flora of Azerbaijan and neighboring countries shows that some species mentioned in the flora of Azerbaijan have already been brought to the subendemic level. These subendemics are also considered Caucasian endemics and are species described for the first time from Azerbaijan (Table 2).

Since some of the sub-endemic species distributed in the Shahdag area have a wide range, their status should be upgraded: *Veronika cristagalli* Stev., *Thesium procumbens* C.A. Mey., *Podospermum canum* C.A. Mey., *Gentianella umbellata* (Bieb.) Holub, *Euphorbia iberica* Boiss., *Cousinia cynaroides* (Bieb.) C.A. Mey., *Cirsium rhizocephalum* C.A. Mey., *Carex caucasica* Stev. and *Cachrys microcarpa* Bieb.

Based on the study of literature sources and our own research, it was determined that 7 species of Azerbaijani endemic and 88 subendemic species are distributed in the territory of Shahdag NP, of which 69 are described only from the Caucasus and are considered endemic to the Caucasus, while others are distributed as far as the borders of Türkiye and Iran (Figs. 1-6).

Relict species are plants preserved in a limited area belonging to the Tertiary and Quarternary periods. In terms of determining the development history of the area flora, the analysis of the relict species and their distribution on different geographical elements was carried out. It was determined that there are 24 relict species in Shahdag flora. In the area, relict species *Mespilus germanica* L., *Parietaria elliptica* C. Koch, *Datisca cannabina* L., *Viola sicheana* W. Beck., *Platanus orientalis* L., *Juglans regia* L., *Rumex scutatus* L., *Krascheninikovia ceratoides* (L.) Gueldents., *Chenopodium foliosum* Aschers., *Phelipanche purpurea* (Jacq.)


Sojak, *Juncus effusus* L., *Thymus nummularius* Bieb., *Rhynchocorys elephas* (L.) Griseb, etc. are found in the composition of various botanical groupings. In general, endemic and relict species in the vegetation of Shahdag area are distributed mainly in meadow vegetation, floodplain and watery meadows, sedge-grass-legume-forb alpine meadows, scree vegetation and shrubby forb vegetation.

There are also some species in the area that have a very narrow range and are in danger of disappearing. Assessment of rare plants according to criteria can be done according to the smallest taxonomic unit. At this time, taking into account the geographic or political area, the condition of populations should be assessed based on the distribution range of the assessed taxon in the wild flora. Each taxon should be assessed against all criteria and assigned an appropriate level of threat. In general, assessment on criteria should be carried out not only by personal expeditions, but also by matching existing scientific materials, clarifying flora and consulting with other specialists. Thus, during the floristic research conducted in the area, the ranges and bioecological characteristics of rare and endangered species were determined, taxa that are few in number, narrowing in ranges and facing the threat of extinction were selected, as well as herbarium materials were collected from different areas of the territory.

In general, 14 species were evaluated as vulnerable (VU), 10 species as Near Threatened (NT), and 5 species as endangered (EN) in the Shahdag area.

The species with special status from the II edition of the Red Book (2013), such as *Dactylorhiza flavescens* (C. Koch) Holub, *Limodorum abortivum* (L.) Sw., *Orchis ustulata* L., *Gladiolus imbricatus* L., *Colchicum speciosum* Stev., *Primula algida* Adams., *Alcea kusariensis* (İljin) İljin, are not included in the III edition of the Red Book (2023), as they are widespread in other area. Currently, the following species in the Shahdag NP have the status of rarity not only in the NP, but also in the biodiversity of Azerbaijan as a whole.

Thus, 23 species identified as rare or endangered in the area under study were evaluated in 3 categories according to IUCN criteria and sub-criteria (Table 3).



Figures 1–6. Endemic and subendemic species of Shahdag National Park (Azerbaijan). Fig. 1: *Angelica sachokiana* (Karjag.) M. Pimen. et V. Tichomirov. Fig. 2: *Astragalus humilis* subsp. *thedodori* (Grossh.) Hashim. Fig. 3: *Campanula meyeriana* Rupr. Fig. 4: *Cirsium macrocephalum* C.A. Mey. Fig. 5: *Betonica nivea* Stev. Fig. 6: *Pseudovesicaria digitata* (C.A. Mey.) Rupr.

Species	IUCN	IUCN	Limiting factors
<i>Woodsia alpina</i> (Bolt) S.F. Gray	NT	In small numbers, in limited areas	Rare species. Anthropogenic factors, sensitivity to environmental factors, very limited range.
<i>Juniperus foetidissima</i> Willd.	NT	Wide distribution areas that are more susceptible to adverse effects.	Rare species. Cattle grazing, fires, unsystematic felling, weak natural regeneration.
<i>Gagea glacialis</i> C. Koch	NT	In small number	Rare species. Decorative plant. In danger of decreasing under the influence of anthropogenic and zoogenic factors.
<i>Herminium monorchis</i> (L.) R. Br.	VU D2	In small number in the only distribution area (Shahdag).	Rare species. Climate change, grazing and trampling.
<i>Himantoglossum formosum</i> (Stev.) C. Koch	EN A2acd+3bc; C1	They are few in number and are in danger of disappearing completely.	Rare, relict species and endemic to the Caucasus. Grazing, trampling and being demanding to a special habitat.
<i>Crocus adami</i> J. Gay	VU B1ab (iii,iv) + 2ab(iii)	A continuous decrease is observed.	Rare species. Decorative plant. Anthropogenic factors, being demanding to a special habitat.
<i>Crocus speciosus</i> Bieb.	VU B1ab(iii) + 2ab(iii)	Decrease of fragmented area quality.	Rare species. Gathering as a decorative, spicy and medicinal plant.
<i>Iris caucasica</i> Stev.	NT	Intensive gathering.	Rare species. Decorative plant. Endemic to the Caucasus. The main limiting factor is anthropogenic influence.
<i>Galanthus caucasicus</i> (Baker) Grossh.	EN B1ab (i,ii,v)c(i,ii,ii) + 2b(i)c	It does not create population.	Rare species. Decorative plant. Endemic to the Caucasus. Gathering as decorative plants, mowing of grass, grazing of cattle.
<i>Aquilegia olympica</i> Boiss.	NT	The population tends to decline.	Rare species. Gathering by the population as an ornamental plant, trampling, grazing and sensitivity to climatic factors
<i>Ranunculus arachnoideus</i> C.A. Mey	VU A2c+3c	Natural reserves are low.	Rare species. Sensitive to environmental factors, short vegetative period, and weak self-regeneration.
<i>Corydalis alpestris</i> C.A. Mey	VU D2	Dependence on anthropogenic influences is observed	Rare species. Anthropogenic factors and climate changes.
<i>Dianthus ruprechtii</i> Schischk	VU D2	The population is small.	Rare species. Endemic to the Caucasus. Habitat is limited, anthropogenic factors, grazing, trampling and mowing.
<i>Gypsophila capitata</i> Bieb.	EN B1ab (iii) + 2ab(iii)	The number of individuals decreases, the distribution area is limited.	Rare species. Endemic to the Caucasus. Climate and anthropogenic factors have caused habitat degradation.
<i>Betula raddeana</i> Trautv.	VU C2a(ii)	A decreasing trend is observed in the population.	Rare species. Endemic to the Caucasus. Decorative plant. Anthropogenic and natural factors are the main limiting factors.

<i>Saxifraga exsarata</i> Vill.	EN B2 ab(i,ii,iii,iv)	Range is narrowing	Rare species. Habitat degradation, weak population regeneration and anthropogenic factors.
<i>Rosa azerbaijdzhanica</i> Novop. et Rzazade	EN B2ab (ii,iii, iv,v)	The habitat is limited, it is completely endangered.	Azerbaijan endemic and rare species. Collection of its flowers and fruits by the population as a medicinal plant.
<i>Astragalus kubensis</i> Grossh.	VU D2	Population decline has accelerated.	Azerbaijan endemic and rare species. The number has decreased due to anthropogenic effects and soil erosion.
<i>Parnassia palustris</i> L.	NT	It tends to decrease.	Rare species. Anthropogenic factors and sensitivity to climatic factors.
<i>Carum caucasicum</i> (Bieb.) Boiss.	NT	It has small populations in limited areas.	Rare species. Sensitivity to adverse climate is observed.
<i>Symphyloloma graveolens</i> C.A.Mey.	NT	The dependence of natural regeneration on the negative effects of climate is observed.	Rare species. Endemic to the Caucasus. Sensitivity to environmental factors, grazing and degradation of distribution areas.
<i>Cladochaeta candidissima</i> (Bieb.)	VU A2c+3c; B1ab(i,iii,iv)	Anthropogenic factors have caused rapid population decline.	Rare species. Endemic to the Caucasus. Spontaneous grazing, medicinal plant, distribution mainly in the flood river beds.
<i>Cynoglossum holosericeum</i> Stev.	VU A2c + 3c	Distributed in limited areas, the population is decreasing due to the influence of changing climate.	Rare species. Endemic to the Caucasus. Medicinal plant. Small number and presence of their ranges in areas with extreme climatic conditions.
<i>Trigonocaryum involucratum</i> (Stev.) Kusn.	VU A2c + 3c	Weak natural regeneration of populations.	Rare species. Anthropogenic factors, floods, and strong erosion processes.
<i>Scrophularia minima</i> Bieb.	NT	Their population is found in small numbers.	Rare species. Endemic to the Caucasus. Anthropogenic factors, distribution in extreme climatic conditions, sensitivity to acute climate change
<i>Veronica minuta</i> C.A. Mey.	NT	Natural recovery in their populations is weak.	Rare species. Endemic to the Caucasus. Since it has a short vegetative period, self-regeneration is weak.
<i>Cetraria islandica</i> (L.) Ach. (= <i>Lichen islandicus</i> L.)	VU A2c +3c	Number is small, the amount and range are about to be exhausted.	Rare species. Used as a medicinal plant in folk medicine since ancient times. Anthropogenic factors (intensive grazing and trampling) and use of mountain meadows.
<i>Usnea florida</i> (L.) F.M. Wigg.	VU D2	Their population is decreasing.	Rare species. It is used as a medicinal plant in folk medicine and medicine. Anthropogenic factors and climatic effects may threaten the extinction of the species.
<i>Lobaria pulmonaria</i> (L.) Hoffm.	VU D2	Their population is decreasing.	Rare species. Medicinal plant. Anthropogenic factors and climatic effects may threaten the extinction of the species.

Table 3. Regional assessment of rare and endangered species of Shahdag area according to IUCN criteria.

There are also some species that are considered rare for Shahdag NP but relatively widespread in other areas: *Aconitum nasutum* Fisch. ex Reichenb., *Pulsatilla albana* (Stev.) Bercht. et Persl., *Pterocarya fraxinifolia* (Poir) Spach. (= *P. pterocarpa* (Michx.) Kunth. ex Iljinsk), *Pyrola rotundifolia* L., *Primula ruprechtii* Kusn., *Viola caucasica* Kolen., *Salix kuznetzowii* Laksch. ex Goerz, *Alchemilla grossheimii* Juz., *Pyracantha coccinea* M.Roem. *Asyneuma campanuloides* (Bieb. ex Sims.) Bornm., *Pyrethrum carneum* Bieb., *Atropa bella-donna* L. (= *A. caucasica* Kreyer). Care should be taken to protect these species as well.

REFERENCES

- Akhundov G.F., 1973. Endemics of the flora of Azerbaijan: / abstr. dis. doctor. biol. Sciences. Baku, 44 pp.
- Asgarov A.M., 2006–2008. Higher plants of Azerbaijan (conspectus of the flora of Azerbaijan). Baku, Elm, Vol. I: 284 pp., Vol. II: 284 pp., Vol. III: 244 pp.
- Asgarov A.M., 2016. The Plant World of Azerbaijan. Baku: Elm, pp. 344–361.
- The Red Book of the Republic of Azerbaijan. Rare and endangered species of plants and mushrooms, 2013. II edition, Baku, 673 pp.
- The Red Book of the Republic of Azerbaijan. Flora, 2023. III edition, Türkiye, 506 pp.
- Babakishiyeva T.J. & İbadullayeva S.J., 2021. Rare plants of Ganja-Kazakh region. Ganja, 225 pp.
- Grosseim A.A., 1939. Analysis of the flora of the Caucasus. AzFAN USSR, 230 pp.
- Hajiyev V.J. & Musayev S.H., 1996. Plants and plant formations recommended for the “Red” and “Green” books of Azerbaijan. Baku, Elm, 40 pp.
- İbadullayeva S. & Mustafayev A.B., 2013. Protection of some rare and critically threatened medicinal plants in the Azerbaijan flora. Journal of Biology and Life Science, 4: 145–152.
<https://doi.org/10.5296/jbls.v4i1.2827>
- İbadullayeva S.J. & Huseynova İ.M., 2021. An Overview of the Plant Diversity of Azerbaijan. Biodiversity, Conservation and Sustainability in Asia, 1: 431–478.
<https://doi.org/10.1007/978-3-030-59928-7>
- İbadullayeva S.J., Mustafayev A.B. & Shiraliyeva G.Sh., 2014. Classification of the vegetation of the high mountainous regions of the Greater Caucasus//Annals of ANAS, Biology series, 3: 58–65.
- IUCN Red List Categories and Criteria: Version 3.1., 2001. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge: UK. ii + 30 pp.
- Grosseim A.A., 1939–1967. Flora of the Caucasus. M.: Nauka, vol. 1–7.
- Musayev S.G., 2005. Revision of endemic species of Azerbaijan. Proceedings of Biological Sciences of ANAS, 1–2: 84–96.
- Mustafayev A., 2019. Taxonomic and Floristic Composition of Flora of Shahdagh National Park (Azerbaijan). Bulletin of Science and Practice, 5: 24–31. <https://doi.org/10.33619/2414-2948/45/03>
- Portenier N.N., 2000. Methodological issues of allocation of geographical elements of the flora of the Caucasus. Botanical Journal, 6: 76–84.
- Tolmachev A.I., 1986. Methods of comparative floristics and problems of florogenesis. Siberian branch, Novosibirsk: Nauka, 195 pp.