

# ***Cistus* ×*skanbergii* Lojac., the ‘phoenix’ hybrid that became extinct in its type locality to revive in the global plant web market**

Salvatore Pasta

Consiglio Nazionale delle Ricerche (CNR), Istituto di Bioscienze e BioRisorse (IBBR), Unit of Palermo, via Ugo La Malfa 153, 90146, Palermo, Italy; e-mail: [salvatore.pasta@ibbr.cnr.it](mailto:salvatore.pasta@ibbr.cnr.it)

## **ABSTRACT**

This paper is focused on the emblematic case story of *Cistus* ×*skanbergii* Lojac., a hybrid rockrose issuing from the breeding between *C. parviflorus* Lam. and *C. monspeliensis* L. To this purpose, a bibliographic research has been carried out to trace the available knowledge about the past and current distribution of this hybrid within its primary distribution range. Extinct in its type locality (Lampedusa Island) and very rare in the Mediterranean Basin, this hybrid proves to be intensely traded worldwide instead. The case of *C. ×skanbergii* shows that the international greening network vehiculated by websites may transform naturally rare plants into cosmopolitan aliens which could become invasive in the near future in several countries subject to Mediterranean or temperate climates. The indiscriminate spread of this rockrose outside its primary distribution range should be discouraged – if not forbidden – also because *C. ×skanbergii* possesses many of the traits (e.g., tolerance to drought and thermal stress, allelopathy, adaptation to fire disturbance) typical to many well-known invasive plants. Additionally, the use of this hybrid rockrose is troublesome because most of the currently traded material has been obtained by crossing several provenances of both parent species under non-natural conditions to obtain the hardiest, hence even more dangerous, breeds.

## **KEY WORDS**

Horticulture; International plant trade; Invasion biology; Mediterranean-type Biomes.

Received 08.09.2023; accepted 22.10.2023; published online 6.11.2023

## **INTRODUCTION**

In one of his contributions to the vascular flora of Lampedusa Island, Lojaccono-Pojero (1885) described *Cistus skanbergii* (Fig. 1) a natural hybrid between *Cistus parviflorus* Lam. and *Cistus monspeliensis* L., two rockrose species which only rarely co-occur throughout the whole Mediterranean area (Fig. 2; Alberto et al., 2020).

The dramatically fast and intense disruption of the island’s territory, due to land overexploitation

during the second half of the 19th century (Pasta & La Mantia, 2003), triggered the fast shrinkage of the local populations of both parent species and of the hybrid itself, originally reported to be extremely common there. As a result, *C. ×skanbergii* was probably extinct in the early 20th century, as reported by Sommer (1908).

This work aims at pointing out the current distribution of *C. ×skanbergii* within its native distribution range, i.e. in the Mediterranean Basin, and highlighting its wide spread due to human activities. Further research was carried out to find out whether

and to what extent this hybrid could be of some interest in the international market of plants traded for ornamental purposes.

## MATERIAL AND METHODS

In order to reconstruct the knowledge on the historical and current distribution of *C. ×skanbergii*, the botanical literature available for Sicily and the Euro-Mediterranean area, as well as several contributions focused on genus *Cistus* (e.g., Rizzotto, 1979; Demoly, 1996, 2006) were consulted. Most data concerning the international trade and the worldwide distribution of the Skånberg rockrose were obtained via *ad hoc* internet searches and/or by using the *iNaturalist* platform (<https://www.inaturalist.org/>).

## RESULTS

### *Ecology and distribution of C. ×skanbergii in its primary distribution range*



Figure 1. Original plate of *Cistus ×skanbergii* (from Lojacono-Pojero, 1885).

Due to its putative hybrid origin, *C. ×skanbergii* should theoretically be present everywhere its parent species co-occur. As matter of fact, this case is not so frequent: according to literature, in fact, this hybrid rockrose has been reported only for western Cyprus (Meikle, 1977; Fig. 2) while the two parent species only seldom co-occur in Crete (Strid, 2016). Despite being closely related to each other from a phylogenetic viewpoint (Civeyrel et al., 2011), the distribution ranges of *C. monspeliensis* and *C. parviflorus* only seldom overlap. The former is mainly a western Mediterranean plant which occurs in few localities in Greece, Cyprus and Crete (Alberto et al., 2020), while the latter is mainly an eastern Mediterranean plant, whose distribution range includes, besides the above-mentioned Island of Lampedusa, Turkey and Turkey-in-Europe, Cyprus, the East Aegean Islands, Greece, Crete, N-Africa (Egypt, Libya and Tunisia). According to the available literature (Böhling et al., 2002; Pignatti et al., 2017–2019) the two species would also have different requirements in terms of soil reaction and light (Table 1).

Demoly (1996) states that *C. ×skanbergii* is mostly sterile in Southern Greece. More recently, Demoly (2006) described *Cistus ×akamantis*, a hybrid between *C. ×skanbergii* and *C. salvifolius* L. living close to the Akamas River in western Cyprus; few years later, the same scholar described a white-flowered form of *C. ×skanbergii* from the same place (Demoly, 2008).

The only claim about the occurrence of *C. parviflorus* elsewhere in Italy refers to Apulia; this record, reported in the first edition of “Flora Europaea” (Tutin et al., 1968), has not been supported neither by herbarium specimens nor by recent observations (Rizzotto, 1979; L. Forte and R.P. Wagensommer, *pers. comm.*). Also the past occurrence of *C. parviflorus* in the garrigues near Caltagirone (southeastern Sicily) needs to be confirmed. Indeed, the only mention for this area is rather ambiguous, as Taranto & Gerbino (1845: 46) quoted “*Cistus complicatus*” (a synonym of *C. parviflorus*) as host of *Cynomorium coccineum* L., while few pages before they report only three species of rockroses growing in the area, i.e. *Cistus salvifolius* L., *C. creticus* L. and *C. monspeliensis* L. As a matter of fact, no specimen of *C. parviflorus* was found in the herbarium of Taranto by Gargano et al. (2018).

Taxon, geographical area and source	L	F	R	N	T	K	S
<i>Cistus monspeliensis</i> L.							
S-Aegean islands, Greece (Böhling et al., 2002)	7	4	8	3	8	3	1
Italy (Pignatti et al., 2017–2019)	10	2	2	2	9	4	0
<i>Cistus parviflorus</i> Lam.							
S-Aegean islands, Greece (Böhling et al., 2002)	8	2	8	n.a.	8	6	1
Italy (Pignatti et al., 2017–2019)	11	2	2	2	9	5	0

Table 1. Ellenberg’s Ecological Indicator Values (EIVs) assigned to *Cistus monspeliensis* and *C. parviflorus* within their native distribution range according to different sources. Standard abbreviations of EIVs: L (= Light), F (= Soil moisture), R (= Soil reaction, pH), N (= Nitrogen soil content), T (= Temperature), K (= Continentality) and S (= Salinity); n.a. = not assessed.



Figure 2. Primary (a: Lampedusa Island, Italy; b: Cyprus) and secondary (c: Republic of South Africa; d: California, US; e: New Zealand; f: Spain) distribution range of *Cistus ×skanbergii* (for further details cf. [https://www.inaturalist.org/observations?locale=it&nelat=45.425957498334846&nelng=180&subview=map&swlat=-80.45301214297774&swlng=-180&taxon\\_id=415891](https://www.inaturalist.org/observations?locale=it&nelat=45.425957498334846&nelng=180&subview=map&swlat=-80.45301214297774&swlng=-180&taxon_id=415891)).

### *An unexpected journey: Skånberg Rockrose “world tour”*

Despite its extinction on Lampedusa Island and its extreme rarity throughout the whole Mediterranean Basin, an automatic query made on the internet (October 2022) using the words ‘*Cistus*’ or ‘rockrose’ + ‘*skanbergii*’ (or ‘*skanbergi*’) revealed

a really unexpected truth. In fact, the total number of results (c. 9200 pages) issuing from this quick search shows that this handsome plant, also known as ‘dwarf pink rockrose’, is massively produced by many nurseries and used for private greening purposes all over Europe (UK, The Netherlands, Germany, Italy, Spain) and worldwide (e.g., USA, Australia and New Zealand). Such reproductive

hybrid individuals have been obtained by crossing on purpose the parent plants from Greece or Cyprus.

*C. ×skanbergii* was introduced as an ornamental plant in the 2000s in some green areas of Valencia (Guillot et al., 2000) and in the roadsides of Morella (Castellón), where it escaped and started to spread at the beginning of the 2010s (Senar Lluch, 2014). Similarly, ten years ago (May 2011) *C. ×skanbergii* was also found growing in an abandoned quarry near Gabbro (Tuscany, Italy; see Lazzeri et al., 2015), where it was probably planted by privates or professionals with no idea - or at least with a weird concept - about what habitat restoration is. No data are available on the destiny of these introduced plants during last decade, but it should better be interpreted as a rare casual alien in Tuscany (V. Lazzeri, *pers. comm.*).

During the last 10–15 years the Skånberg rockrose was introduced worldwide in all the countries with Mediterranean-type climate like Chile, Southern Australia, as well as in South Africa and California, where it starts to behave as a casual or fully naturalized alien plant (Fig. 2).

## DISCUSSION

### **Nemo invasor in patria? Evolutionary and ecological drivers**

The evergrowing bulk of knowledge on the most effective biology predictors of invasiveness (Hayes & Barry, 2008) allows to make some more considerations on the specific case of *C. ×skanbergii*.

First, narrow-ranged does not mean harmless (Pasta, 2022). In fact, vascular plants with no close relatives may pose serious problems to the ecosystems of newly colonized areas (e.g., Callaway & Aschenhoug, 2000; Colautti et al., 2017). The dreadful weapon represented by ‘phylogenetic originality’ has been confirmed by many recent studies carried out worldwide, emphasizing that the more distinct are local floras, the more they are subject to be outcompeted by alien invaders (Harvey et al., 2012).

Second, if we consider ecological pre-adaptation (Colautti & Barrett, 2013), *C. ×skanbergii* looks just predestined to spread all over the Mediterranean biomes worldwide. In fact, thanks to their

high stress and disturbance-tolerance (Troia & Laguna Lumbreras, 2015) rockroses can survive and spread in all the arid and fire-prone habitats.

Although homoploid hybrids are rare in nature, nonetheless they exist, as shown by Kadereit (2015). Even if *Cistus* species are with no exceptions diploid (Totta et al., 2017), they may be easily crossed and form plenty of hybrids and intermediates either in nature or under cultivation; for this reason, hybridization has been hypothesized as the major mode of evolution in *Cistus* since the early 20th century (Dansereau, 1940; Demoly, 1996).

The possibility that *C. ×skanbergii* may be able not only to overcome sterility but also to show improved fitness like other *Cistus* hybrids (Navarro-Cano et al., 2017) sheds light on the potential threat posed by Skånberg rockrose. Additionally, hybrid origin represents another favourable trait shared by many invasive plants, as already pointed out by Stebbins (1985) and recently confirmed for many evolutionarily distant plants (e.g., Abbott et al., 2003; Schierenbeck & Ellstrand, 2009; Yakimowski & Rieseberg, 2014).

Moreover, the specific case of this ‘home-made hybrid rockrose’ is particularly troublesome because of its production in non-natural conditions by crossing several provenances of both parent species. In fact, the enhanced genetic diversity issuing from admixture may significantly increase the ecological plasticity and fitness of many introduced plants and animals (Lavergne & Molofsky, 2007).

### **Is it just a question of time (and speed)?**

Mankind started changing the worlds’ floristic assemblages already many thousand years ago, when the first seafarers brought with them the propagules (seeds, tubers, bulbs, etc.) of the food and medicine plants they needed (Merlin, 2003; Fuller et al., 2011; Hofman & Rick, 2018), and the same did for centuries sailors, monks and nomad shepherds following ancient trade routes and carrying with them useful plants to be cultivated on islands after crossing immense oceans or grown in the oases after weeks of wandering across hostile deserts. If compared with past times, the international commerce of potentially dangerous weeds has dramatically increased in terms of speed, inten-

Wide climatic niche (able to thrive under cooler conditions and/or to enjoy global warming)	NE
Disturbance-tolerant (e.g. volcanic eruptions, wildfires, floods, rockfalls)	Y
Hybrid/Polyploid	Y
Dominant species	N
Early reproductive	Y
Long-lasting flowering/fruitletting season	Y
Wide edaphic niche (indifferent to soil chemistry)	NE
Multiple pollination strategy and wide pollination range	NE
Multiple dispersal strategy and wide seed dispersal range	N
Able to produce vegetative propagules	N
Habitat transformer/accumulator (e.g. nitrogen-fixing, allelopathy)	Y
Stress-tolerant (e.g., CAM, C4, salt-tolerant, pollution-tolerant, drought-tolerant)	Y

Table 2. Ecological traits of *C. ×skanbergii* which could favour (Y) or prevent (N) its invasive behaviour and spread outside its primary range. NE = not evaluated, to be checked outside its native range.

sity, frequency and species-richness. Seed and plant delivery via airplanes and online shopping, often used for the illegal commerce of very endangered and internationally protected plants (Vaglica et al., 2017), represents at the same time the worst enemy of rare plants and the best ally for invasive ornamental ones (Humair et al., 2015).

The fact that *C. ×skanbergii* did not turn invasive yet does not mean that it cannot become so in the future, especially if it turns to be able to overcome sterility. Indeed, we should beware of apparent unharfulness (Zenni & Nuñez, 2013), considering that many introduced alien species show a very long lag time before becoming pests (Crooks & Soulé, 1999; Groves, 2006; Lambdon et al., 2008). A synthesis on the ecological traits making *C. ×skanbergii* a ‘dangerous trade item’ is provided in Table 2.

## CONCLUSIONS

How to translate scientific discoveries on biological invasions into everyday’s conservation policy and practice? Perhaps we should start from raising awareness and changing the way of perceiving invasive organisms. In fact, teaching stakeholders and common people to go beyond the aesthetical look of alien plants and to see their impact on natural and semi-natural ecosystems (Andreu et al., 2009; Lindemann-Matthies, 2016; Dehnen-Schmutz & Conroy, 2018) could make them the first detectors of invasion events (see Encarnação et al., 2021, and reference therein). With no doubt, it is more cost-effective to prevent and forbid the intentional introduction of potentially invasive alien plants instead of fighting the very widespread and well-established ones (Hulme, 2003; Hulme et al., 2008).

Perfectly aware of their past and present responsibilities and duties (Dehnen-Schmutz, 2011; Hulme; 2011; Dehnen-Schmutz & Conroy, 2018), the European horticulturalists and the botanic gardens are networking to find out joint policies focused on the most appropriate way of dealing with (and avoiding) invasive ornamental species. Such increasingly tight collaboration between private and public gardening organisations and public research institutions led to the recent publishing of various international codes or guidelines (Heywood & Brunel, 2011), whose authoritative suggestions, however, are still waiting to be taken seriously in many countries, where the introduction of apparently unharful alien plant species such as *C. ×skanbergii* has been encouraged in rural and urban ecosystems until recent times (Nicholls et al., 2001; Fraga i Arguimbau, 2009; Canales-Ide et al., 2022).

## ACKNOWLEDGEMENTS

I thank Angelo Troia and Riccardo Guarino (Department STeBiCeF, University of Palermo, Italy) and the anonymous reviewers, whose comments improved the manuscript, Valerio Lazzeri (Museo di Storia Naturale del Mediterraneo, Leghorn, Italy) for sharing information on the escaped individuals of *C. ×skanbergii* in Tuscany, Jean-Marc Tison (Heyrieux, France) and Frédéric Médail (IMBE, University of Aix-Marseille, France) for helping to

consult the publications of Mr. J.-P. Demoly, Ralf Jahn (Großschirma, Germany) for providing information on the current distribution of the rockroses in Greece, Crete and the Aegean Islands, Luigi Forte (Botanic Museum and Garden of the University of Bari, Italy) and Robert Philipp Wagensommer (Department of Biology, University of Bari, Italy) for their advice about the past presence of *C. parviflorus* in Apulia.

## REFERENCES

- Abbott R.J., James J.K., Milne R.I. & Gillies A.C.M., 2003. Plant introductions, hybridization and gene flow. *Philosophical Transactions of the Royal Society B (Biological Sciences)*, 358 (1434): 1123–1132. <https://doi.org/10.1098/rstb.2003.1289>
- Alberto J., Coello A.J., Fernández-Mazuecos M., García-Verdugo C. & Vargas P., 2020. Phylogeographic sampling guided by species distribution modelling reveals the Quaternary history of the Mediterranean-Canarian *Cistus monspeliensis* (Cistaceae). *Journal of Systematic and Evolutionary*, 59: 262–277. <https://doi.org/10.1111/jse.12570>
- Andreu J., Vilà M. & Hulme P.E., 2009. An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management*, 43: 1244–1255. <https://doi.org/10.1007/s00267-009-9280-1>
- Böhlng N., Greuter W. & Raus Th., 2002. Zeigerwerte der Gefäßpflanzen der Südägäis (Griechenland) - Indicator values of the vascular plants in the Southern Aegean (Greece). *Braun-Blanquetia*, 32: 1–108.
- Callaway R.M. & Aschehoug E.T., 2000. Invasive plants versus their new and old neighbors: a mechanism for exotic invasion. *Science*, 290: 521–523. <https://doi.org/10.1126/science.290.5491.521>
- Canales-Ide F., Zubelzu S., Segovia-Cardozo D. & Rodríguez-Sinobas L., 2022. Assessing the performance of irrigation systems in large scale urban parks: Application to the case of Valdebebas, Madrid (Spain). *Remote Sensing*, 14: 1060. <https://doi.org/10.3390/rs14051060>
- Civeyrel L., Leclercq J., Demoly J.-P., Agnan Y., Quèbre N., Péliissier C. & Otto T., 2011. Molecular systematics, character evolution, and pollen morphology of *Cistus* and *Halimium* (Cistaceae). *Plant Systematics and Evolution*, 295: 23–54. <https://doi.org/10.1007/s00606-011-0458-7>
- Colautti R.I., Alexander J.M., Dlugosch K.M., Keller S.R. & Sultan S.E., 2017. Invasions and extinctions through the looking glass of evolutionary ecology. *Philosophical Transactions of the Royal Society B (Biological Sciences)*, 372: 20160031. <https://doi.org/10.1098/rstb.2016.0031>
- Colautti R.I. & Barrett S.C.H., 2013. Rapid adaptation to climate facilitates range expansion of an invasive plant. *Science*, 342: 364–366. <https://doi.org/10.1126/science.1242121>
- Crooks J.A. & Soulé M.E. 1999. Lag times in population explosions of invasive species: causes and implications. In: Sandlund O.T., Schei P. & Viken Å. (Eds), *Invasive species and biodiversity management*. Kluwer Academic Publishers, Dordrecht, pp. 103–125.
- Dansereau P., 1940. Études sur les hybrides de *Cistes*. *Annales des épiphyties*, sér. 2, 6: 7–26.
- Dehnen-Schmutz K., 2011. Determining non-invasiveness in ornamental plants to build green lists. *Journal of Applied Ecology*, 48: 1374–1380. <https://doi.org/10.1111/j.1365-2664.2011.02061.x>
- Dehnen-Schmutz K. & Conroy J., 2018. Working with gardeners to identify potential invasive ornamental garden plants: testing a citizen science approach. *Biological Invasions*, 20: 3069–3077. <https://doi.org/10.1007/s10530-018-1759-3>
- Demoly J.-P., 1996. Les hybrides binaires rares du genre *Cistus* L. (Cistaceae). *Anales del Jardín Botánico de Madrid*, 54: 241–254.
- Demoly J.-P., 2006. Les hybrides ternaires du genre *Cistus* L. (Cistaceae). *Biocosme Méditerranéen*, 23: 1–15.
- Demoly J.-P., 2008. Nouveautés nomenclaturales pour le genre *Cistus*. *Bulletin de l'Association des Parcs Botaniques de France*, 45: 47–48.
- Encarnação J., Teodósio M.A. & Morais P., 2021. Citizen science and biological invasions: A review. *Frontiers in Environmental Science*, 8: 602980. <https://doi.org/10.3389/fenvs.2020.602980>
- Fraga i Arguimbau P., 2009. Jardineria mediterranea sin especies invasoras. *Generalitat Valenciana, Conselleria de Medio Ambiente, Agua, Urbanismo y Vivienda*, Valencia, 208 pp.
- Gargano M.L., Domina G. & Venturella G., 2018. The neglected herbarium of Emanuele Taranto Rosso (Sicily, 1801–1887). *Atti della Società Toscana di Scienze Naturali, Memorie, serie B*, 125: 1–7. <https://doi.org/10.2424/ASTSN.M.2018.10>
- Groves R.H., 2006. Are some weeds sleeping? Some concepts and reasons. *Euphytica*, 148: 111–120.
- Guillot Ortiz D., Mateo Sanz G. & Rosselló Picornell J.A., 2009. Claves para la flora ornamental de la Provincia de Valencia. *Monografías de Bouteloua*, 1: 1–320 pp.
- Harvey K.J., Nipperess D.A., Britton D.R. & Hughes L., 2012. Australian family ties: does a lack of relatives help invasive plants escape natural enemies? *Biological Invasions*, 14: 2423–2434. <https://doi.org/10.1007/s10530-012-0239-4>
- Hayes K.R. & Barry S.C., 2008. Are there any consistent

- predictors of invasion success? *Biological Invasions*, 10: 483–506.  
<https://doi.org/10.1007/s10530-007-9146-5>
- Heywood V.H. & Brunel S., 2011. Code of Conduct on Horticulture and Invasive Alien Plants. Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention). Council of Europe Publishing, Nature and Environment, Strasbourg, no. 162, 93 pp.
- Hulme P.E., 2003. Biological invasions: winning the science battles but losing the conservation war? *Oryx*, 37: 178–193.  
<https://doi.org/10.1017/S003060530300036X>
- Hulme P.E., 2011. Addressing the threat to biodiversity from botanic gardens. *Trends in Ecology & Evolution*, 26: 168–174.  
<https://doi.org/10.1016/j.tree.2011.01.005>
- Hulme P.E., Bacher S., Kenis M., Klotz S., Kuhn I., Minchin D., Nentwig W., Olenin S., Panov V., Pergl J., Pyšek P., Roques A., Sol D., Solarz W. & Vilà M., 2008. Grasping at the routes of biological invasions: a framework to better integrate pathways into policy. *Journal of Applied Ecology*, 45: 403–414.  
<https://doi.org/10.1111/j.1365-2664.2007.01442.x>
- Humair F., Humair L., Kuhn F. & Kueffer C., 2015. E-commerce trade in invasive plants. *Conservation Biology*, 29: 1658–1665.  
<https://doi.org/10.1111/cobi.12579>
- Lavergne S. & Molofsky J., 2017. Increased genetic variation and evolutionary potential drive the success of an invasive grass. *Proceedings of the National Academy of Sciences of the United States of America*, 104: 3883–3888.  
<https://doi.org/10.1073/pnas.0607324104>
- Lazzeri V., Sammartino F., Campus G., Caredda A., Mascia F., Mazzoncini V., Testa N. & Gestri G., 2015. Note Floristiche Tosco-Sarde II. Novità regionali e locali e considerazioni tassonomiche per le regioni Sardegna e Toscana. *Annali del Museo Civico di Rovereto, Sezione: Archeologia, Storia, Scienze Naturali*, 30 (2014): 331–368.
- Lindemann-Matthies P., 2016. Beasts or beauties? Laypersons’ perception of invasive alien plant species in Switzerland and attitudes towards their management. *NeoBiota*, 29: 15–33.  
<https://doi.org/10.3897/neobiota.29.5786>
- Lojacono-Pojero M., 1885. Una escursione botanica in Lampedusa. *Il Naturalista siciliano*, 4: 92–96.
- Meikle R.D., 1977. *Flora of Cyprus*. Vol. 1. Bentham Moxton Trust, Royal Botanic Gardens, 832 pp.
- Navarro-Cano J.A., Schwienbacher E., Sánchez-Balibrea J. & Erschbamer B., 2017. The role of seed traits as segregation factors of hybrids in wild populations of *Cistus* (Cistaceae). *Plant Biosystems*, 151: 530–538.  
<https://doi.org/10.1080/11263504.2016.1186125>
- Nicholls C.I., Parrella M. & Altieri M.A., 2001. The effects of a vegetational corridor on the abundance and dispersal of insect biodiversity within a northern California organic vineyard. *Landscape Ecology*, 16: 133–146, 2001.  
<https://doi.org/10.1023/A:1011128222867>
- Pasta S., 2022. Never underestimate Sicilians: some case histories dealing with narrow native ranges, deliberate introduction, claimed extinction and predictable plant invasion worldwide. *Flora Mediterranea*, 32: 403–420. <https://doi.org/10.7320/FIMedit32.403>
- Pasta S. & La Mantia T., 2003. Note sul paesaggio vegetale delle isole minori circumsiciliane. II. La vegetazione pre-forestale e forestale nelle isole del Canale di Sicilia. *Annali Accademia Italiana di Scienze Forestali*, 51: 77–124.
- Pignatti S., Guarino R. & La Rosa M. (Eds.), 2017–2019. *Flora d’Italia*. 2a edizione. Edagricole-New Business Media, Bologna - Milano, 4 volumes.
- Rizzotto M., 1979. Ricerche tassonomiche e corologiche sulle Cistaceae. 1. Il genere *Cistus* L. in Italia. *Webbia*, 33: 343–378.  
<https://doi.org/10.1080/00837792.1979.10670127>
- Senar Lluch R., 2014. Aportaciones de la distribución de la flora de la Provincia de Castellón II. *Flora Montiberica*, 57: 3–16.
- Sommier S., 1908. *Le Isole Pelagie Lampedusa, Linosa, Lampione, e la loro Flora*. Con un elenco completo delle piante di Pantelleria. Stabilimento Pellas, L. Chiti e Successori, Firenze, 345 pp.
- Strid A., 2016. *Atlas of the Aegean Flora*. Englera, 33, Part 2 (Maps). Botanic Garden and Botanical Museum Berlin, Freie Universität Berlin, Berlin.
- Taranto (Rosso) E. & Gerbino X., 1845. *Catalogus Plantarum in Agro Calata-Hieronensi Collectarum*. Fasciculus I. Catanæ: ex Typographia Josephi Musumeci-Papale, 1–50 pp. (+ 1 “Errata Corrige”).
- Totta C., Rosato M., Ferrer-Gallego P., Lucchese F. & Rosselló J.A., 2017. Temporal frames of 45S rDNA site-number variation in diploid plant lineages: Lessons from the rock rose genus *Cistus* (Cistaceae). *Biological Journal of the Linnean Society*, 120: 626–636.  
<https://doi.org/10.1111/bij.12909>
- Troia A. & Laguna Lumbreras E., 2015. On the pyrophytism in the Mediterranean area. *Journal of Arid Environments*, 120: 1–3.  
<https://doi.org/10.1016/j.jaridenv.2015.04.003>
- Tutin T.G., Heywood V.H., Burges N.A., Valentine D.H., Walters S.M. & Webb D.A. (Eds.), 1968. *Flora Europaea*. Cambridge University Press, Cambridge, London, New York, Melbourne. Vol. 2: Rosaceae to Umbelliferae: xxvii + 475 pp.
- Vaglica V., Sajevo M., McGough H.N., Hutchison D., Russo C., Gordon A.D., Ramarosandratana A.V.,

Stuppy W. & Smith M.J., 2017. Monitoring internet trade to inform species conservation actions. *Endangered Species Research*, 32: 223–235.  
<https://doi.org/10.3354/esr00803>

Zenni R.D. & Nuñez M.A., 2013. The elephant in the room: the role of failed invasions in understanding invasion biology. *Oikos*, 122: 801–815.  
<https://doi.org/10.1111/j.1600-0706.2012.00254.x>