

Diversity of the land snail fauna of Llogara National Park (Albania) focusing on the distribution area of *Chondrula lugorensis* A.J.Wagner, 1914 (Gastropoda Enidae)

Ina Nasto¹, Denada Sota¹, Brunilda Veshaj¹, Geolind Çobaj¹, Encora Çelohoxhaj², Marsel Javori¹, Nexhip Hysolako² & Emirjeta Adhami³

¹Department of Biology, Faculty of Technical and Natural Sciences, University Ismail Qemali Vlore, Albania

²Regional Administration of Protected Areas, Vlore, Albania

³Institute for Nature Conservation in Albania (INCA)

*Corresponding author: e-mail: ina.nasto@yahoo.com

ABSTRACT

Regular surveys of biodiversity are important for monitoring changes in the richness and species composition of mollusk fauna, especially in mountainous areas and fragmented and isolated habitats that are biodiversity hotspots, characterized by endemic species with isolated spread. This paper presents a list of the terrestrial mollusk fauna of Llogara National Park (Albania) and includes data on the distribution area of the endemic mollusk *Chondrula (Chondrula) lugorensis* Wagner, 1914. Snails and slugs were collected using visual searches and leaf litter collections during the spring and autumn seasons of 2020 and 2021. A total of 52 terrestrial gastropods were identified, representing 13 species from nine families. The study also sampled 24 individuals, both alive and deceased, of the endemic species *Chondrula (Chondrula) lugorensis* (Gastropoda Enidae), which allowed for creating a distribution map and identifying its habitat.

KEY WORDS

Chondrula lugorensis; malacofauna; Llogara; endemic; land snail; Albania.

Received 22.05.2023; accepted 26.10.2023; published online 30.12.2023

INTRODUCTION

Terrestrial mollusks represent one of the most speciose groups of terrestrial animals on Earth, with about 28,000 recognized species (MolluscaBase 2022) and an estimated number of undescribed taxa ranging between 11,000 and 40,000 (Lydeard et al., 2004; Rosenberg, 2014; Goudeli et al., 2021). The heterogeneous structure of the habitat and the geological fragmentation of the Balkan peninsula make it one of the major hotspots of terrestrial mollusk diversity in the Holarctic region (Fehér & Eröss, 2009; Neubert et al., 2019). Albania remains among the countries with the least data regarding terrestrial

mollusk fauna, probably due to the geopolitical situation and the difficult accessibility of this country (Fehér & Eröss, 2009).

The chronological reconstruction of studies of terrestrial mollusk fauna in Albania dates to the first expeditions in the mid-19th century by SCHLÄFLI, Westerlund, Blanc, and Von Martens (Mousson, 1859; Martens, 1889; Westerlund & Blanc, 1879; Fehér & Eröss, 2009). In the early 20th century, a number of data from Apfelbeck, Buljubasić, Führer, Klapotocz, Petrović, Sturany, Wagner, Winneguth, and Wohlberedt were collected and presented in two monographs (Wohlberedt, 1909; Sturany & Wagner, 1915), followed by other publications during

the First and Second World Wars (Csiki, 1923; Kümmerle, 1926; Wagner, 1913; Wagner, 1919). Then followed by a limited number of studies during the period of the communist regime mainly published by the Academy of Sciences of Albania and the University of Shkodra, (Jaekel et al., 1957; Jaekel & Schmidt; 1961; Dhora, 1976; 1978a, b; 1981; 1983; 1985; 1988a, b; 1990).

After the political transition, the explorations of terrestrial mollusks fauna started again with some expeditions organized in collaboration between Albanian and Austrian researchers (Welter-Schultes 1996; Riedel & Welter-Schultes, 1996; Dhora & Welter-Schultes, 1996; 1999a; b). Until 1996, the fauna of terrestrial mollusks counted 237 species of terrestrial and freshwater mollusks for Albania (Dhora, 1994; 1995; 1995–96; 1997a; 1997b; 1997–2000; 1999a; 1999b; Dhora & Welter-Schultes, 1996). Until this moment 48 taxa have been described, among which some new species for the science have been reported (Subai 1997, 1999a, b; Dhora & Welter-Schultes, 1999a, b; Eröss et al., 1999, 2005, 2006; Riedel et al., 1999; Fehér et al., 2001; Fehér 2004; Fehér & Drimmer 2004; Subai & Fehér, 2006; Fehér & Eröss, 2009; Lika et al., 2021).

Therefore, in addition, we are presenting some data about the biodiversity of terrestrial mollusks of Llogara National Park. This paper also tries to re-present the distribution areal of the species *Chondrula lugorensis* Wagner, 1914 in the Llogara national park, described for the first time by Sturany & Wagner (1915) and recently reported by Dhora & Welter-Schultes (1996), Fehér & Eröss (2009) and Welter-Schultes (2012).

MATERIAL AND METHODS

Study area

Llogara National Park is located in the southwest region of Albania, along the Ceraunian Mountain range that stretches along the Ionian Sea. The Karaburun peninsula is located on the northeastern side of the Strait of Otranto, where the Adriatic Sea meets the Ionian Sea. While the floristic richness of the Karaburun-Çika Mts Protected Area has not been fully documented, this complex ecosystem is considered to be the most diverse part of Albania in

terms of species with conservation interest. The National Park is home to nine endemic vascular plants, such as *Arenaria cikaea*, *Edraianthus caespitosus*, and *Dianthus sylvestris subsp. alboroseus*, *Hypericum haplophyloides*, *Limonium himariense*, *Nocca cikaea*, *Reichardia albanica*, *Sesleria albanica*, and *Viola acrocerauniensis*. Additionally, the Llogara National Park is home to four endemic terrestrial mollusks, such as *Chondrula lugorensis*, *Allaegopsis amphikypellon*, *Monacha emigrata senitshika*, and *Helicigona fuchsia*, as well as one endemic carabid, *Dorcadion albanicum*.

The Llogara National Park is also home to several globally threatened species, including the Horse Chestnut (*Aesculus hippocastanum*) and Autumn Snowdrop (*Galanthus reginae-olgae*). Six other species, namely *Fritillaria graeca*, *Anacamptis pyramidalis*, *Ruscus aculeatus*, *Ophrys helenae*, *Ophrys oestrifera*, and *Ramonda serbica*, are of great conservation interest due to their protection under the Habitat Directive (92/43/EEC) and Berne Convention.

The Park is also home to 52 plant species that are part of Albania's red list of protected flora, but the complete number of flora and fauna species is not yet fully known. The plant cover vegetation in the Llogara National Park is dominated by various habitat types, including vegetated sea cliffs of the Mediterranean coasts with endemic *Limonium* spp., stable xerothermophilous formations with *Buxus sempervirens* on rock slopes, Hellenic beech forests with *Abiesborisii-regis*, *Quercus ilex*, and *Quercus rotundifolia* forests, *Quercus macrolepis* forests, and High Oro-Mediterranean pine forests.

In addition to the diverse plant life, the LNP is also home to a rich variety of invertebrate and vertebrate species. The area is known to house 98 fish species (both freshwater and marine), 10 amphibian species, 28 species of reptiles, 105 species of birds (mostly in Orikumi lagoon), and 55 mammal species (Final Report on the Study on Landscape of Llogara National Park in Albania).

Survey methods

The surveys were conducted in the Llogara National Park area, including Cika Mountain and the Palasa River (Fig. 1), which contain native and non-native snails, both macro- and micro-mollusks, in a variety of habitats. Sampling was carried out from



Figure 1. Map of the areas visited during the expeditions. 1. The path of Caesar; 2. Cika Maintain; 3. Panorama; 4. Palasa torrent.

Latitude	Longitude	Number of specimens
40° 12.950'N	19° 36.380'E	2
40° 12.830'N	19° 36.659'E	3
40° 11.911'N	19° 35.498'E	1
40° 12.941'N	19° 36.421'E	1
40° 12.967'N	19° 36.554'E	3
40° 12.895'N	19° 36.474'E	2
40° 12.773'N	19° 36.705'E	1
40° 12.648'N	19° 36.945'E	2
40° 12.505'N	19° 37.608'E	3
40° 12.229'N	19° 35.529'E	1 (empty shell)
May be transported by water		
40° 11.956'N	19° 35.389'E	1 (empty shell)
May be transported by water		

Table 1. Location of sampled specimens (Llogara National Park, Albania).

February to September in 2020, and 2021. In each area, five sampling sites of 10 m² were selected, spaced at least 100 m from each other along a transect to maximize the number of different sampled microhabitats. The Cika mountain sites were visited three times as species had been previously reported in that area, while other sites were visited only once to confirm the presence or absence of the species. However, during two visits to the Cika mountain area and two other areas, *Chondrula lugorensis* was found (Table 1, Fig. 2).

Sampling began with a visual search of suitable microhabitats, including plants, by expert snail gatherers lasting 15 minutes with five people at each site. This was followed by collecting individuals at random within each site, including places with deep soil accumulation and around the bases of rocks and trees. All collected material was stored in 70% alcohol and taken to the laboratory for further processing. The samples collected using both methods were identified at the lowest possible taxonomic level, and in some cases, specimens were referred only to a family, subfamily, or genus. Any species readily identified in the field were simply counted.

The research was aided by students from the University “Ismail Qemali” Vlore, who were prepared for all stages of sampling, identification, and data collection on the species. Although it was often difficult to determine whether an individual died prior to collection or during storage, all samples were processed, and live and dead (shell only) counts were pooled, except for *Chondrula lugorensis*, which counts were divided into living and dead specimens. A subset of live specimens was stored in 95% ethanol as representative vouchers for future analysis. All vouchers have been cataloged and deposited in the “Ismail Qemali” University in Vlore.

RESULTS

During the four expeditions, 13 species belonging to nine different taxonomic groups were collected and identified. All specimens have been identified at the species level, with the exception of the genus *Montenegrina* which will be examined much more carefully before declaring the species (Table 2).

Most of the species were alive at the time of sampling, with the exception of some individuals, who preferred to sample only the shell for a minor

Species	The Path of Caesar	Cika Maintain	Palasa torrent	Panorama
<i>Cochlostoma auritum</i> (Rossmässler, 1837)	+	+		+
<i>Cochlostoma tessellatum</i> (Rossmässler, 1837)	+	+	+	+
<i>Pomatias elegans</i> (O.F. Müller, 1774)	+	+		+
<i>Herilla illyrica</i> (Möllendorff, 1899)			+	
<i>Montenegrina</i> sp. O. Boettger, 1877		+		+
<i>Cernuella virgata</i> (Da Costa, 1778)		+		
<i>Helicodonta obvoluta</i> (O.F. Müller, 1774)	+		+	
<i>Aegopis acies</i> (Férussac, 1832)			+	+
<i>Aegopis septentrionalis laughofferi</i> A.J. Wagner, 1914	+			+
<i>Chondrula lugorensis</i> A.J. Wagner, 1914		+		
<i>Chondrula microtragus</i> (Rossmässler, 1839)		+	+	
<i>Lauria cylindracea</i> (Da Costa, 1778)	+		+	
<i>Radix peregra</i> (O.F. Müller, 1774)	+		+	

Table 1. Mollusca fauna was identified in Llogara National Park in four different locations: 1. The path of Caesar; 2. Cika Maintain; 3. Panorama; 4. Palasa torrent.

impact on the population. All families are represented with a maximum of 2 species. The families with the greatest number of species are Zonitidae (2 species), Enidae (2), Cochlostomatidae (2), and Clausiliidae (2).

Data on *Chondrula lugorensis* A.J. Wagner, 1914 distribution

Chondrula (Chondrula) lugorensis, is a species of terrestrial Gastropoda in the family Enidae. First, this species was described in 1915 in an article published in the Memoranda of the Imperial Academy of Sciences, Mathematical and Scientific Class, where some of the terrestrial mollusk species found in Albania and its surroundings were described and reported (Sturany & Wagner, 1915).

According to Sturany and Wagner (1915) *C. lugorensis* is a small gastropod, 8–11 cm long and 3.5–4 cm large with a dirty yellowish-grey shell, translucent, yellowish opaque near the aperture, coarsely and irregularly striated, 8 weakly convex whorls, last whorl not descending before aperture, aperture simple without teeth, margin straight and thin, with a white lip inside, sometimes forming some teeth-like structures, margins connected at parietal side by a thin layer, but do not approach each other much, umbilicus very narrow (Fig. 2).

Later on, *C. lugorensis* was reported in several articles but in the same area of distribution (Dhora, 2002 a and b, 2004; Fehér & Eröss, 2009 a and b; Welter-Schultes, 2012).

The actual distribution of *C. lugorensis* known since now are:

Qafa e Llogarait (South Albanian coast)
Periferi Vlorë, Mali i Çikës, 2 km W
Maja e Qorrës along the ridge to Maja e Çikës,
1840 m a.s.l.,
Periferi Tepelene, ca. 1 km SE
Maja e Klogjurit, 1779 m a.s.l.,

Thanks to an assessment done in 2010 on the distribution area of this species, *C. lugorensis* was declared by the IUCN as a Near Threatened species according to Feher (2013). According to IUCN, the number of currently known locations is less than five, and the known area of occupancy (AOO) is less than 20 km² of occurrence (EOO), the number of locations, the number of subpopulations or the number of mature individuals are declining or extremely fluctuating. There is no major threat to this species; therefore, it is assessed as Near Threatened (NT), almost qualifying for Vulnerable D2. During the monitoring and sampling activities, about 20 individuals of *C. lugorensis* were sampled in four sites in the Cika mountain area (Fig. 3).



Figure 2. *Chondrula (Chondrula) lugorensis* Wagner, 1914 (photo by I. Nasto).




Figure 3. Map of the actual known distribution of (*C. lugorensis* Feher, (2013), ascertained from the sampling campaigns during 2019–2021 (Map created by Ronny Dobbelssteijn).

Most of the individuals sampled were chosen dead, due to the limitation of the impact on the population of the species. Live individuals of different sizes were sampled in order to clearly describe the species. The species was often accompanied by *Cochlostoma tessellatum* (Rossmässler, 1837) and *Herilla* sp.

Habitat and ecology

Currently, we know little about the ecology and habitat of *C. lugorensis* but knowing its distribution we can give some important characteristics of the habitat. *Chondrula lugorensis* is a typical mollusk associated with a karstic habitat. The distribution of the species is the same as it was evaluated in 2010. The distribution is compatible with the habitat of high Mediterranean pine forest but in the areas of rare forest, characterized by the vegetation of *Sesleria albanica*, *Festuca* sp., and other perennial Poaceae grasses. The species is usually located in humid areas consisting of limestone covered with moss, proof that the habitat is humid most of the time. So far it has been sampled only in limestone areas exposed to the northeast in order to maintain a constant temperature and humidity. This mollusk is often positioned vertically under the rocks of the limestone walls quite sheltered from the wind and sun.

Little is known of its biology, including breeding biology. There is a suggestion that the species breed throughout the year when conditions are suitable.

Often the species hides between roots and moss, and being the same color as the surrounding vegetation, it gives it the opportunity to blend in perfectly with the surrounding environment. In fact, no traces of consumption by animals such as insects or birds have been found. All the individuals found, both alive and dead, had the entire shell, not damaged by possible carnivores. Three damaged shells were found which were rated as very old and damaged by the decomposition process.

ACKNOWLEDGMENTS

All monitoring campaigns in the Llogara National Park were funded through the project “From Mountain to the Sea” Project reference: CEPF–108695,

coordinated by the Institute for Nature Conservation in Albania (INCA). Ethical clearance: we hereby declare that the manuscript is an original work produced by genuine research, never previously published. No data was fabricated or manipulated (including images) to support your conclusions. The present study followed international, national, and institutional guidelines for animal treatment and complied with relevant legislation “for sampling and research on marine protected areas in Albania”. No experiment against Ethical clearance for the use of animals in scientific research has been carried out.

REFERENCES

- Csiki E., 1923. Előszó. In: A Magyar Tudományos Akadémia Balkán-kutatásainak tudományos eredményei. Teleki P. & Csiki E. (Eds.), Vol. 1. Csiki E. állattani kutatásai Albániában - Explorationes Zoologicae ab E. Csiki in Albania Peractae. Pars 1., The Hungarian Academy of Sciences, Budapest. Denkschriften der kaiserlichen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse, 91: 19–138.
- Dhora Dh. & Welter-Schultes F.W., 1996. List of species and atlas of the non-marine molluscs of Albania. Schriften zur Malakozoologie, 9: 90–197.
- Dhora Dh. & Welter-Schultes F.W., 1999a. Mollusc cenoses from different environments in Albania. Schriften zur Malakozoologie, 13: 61–66.
- Dhora Dh. & Welter-Schultes F.W., 1999b. Clausiliidae of Albania: recent finds and biogeographical evaluations (Gastropoda, Pulmonata). Biologia Gallo-hellenica, 25: 15–24.
- Dhora Dh., 1976. Të dhëna nga studimi i molusqeve të Shqipërisë së Ulët Bregdetare. Buletini i Shkencave të Natyrës, 1: 69–72.
- Dhora Dh., 1978. Mbi disa dyflegërorë më të njohur të zonave të Shëngjinit dhe të Vlorës. Buletini i Shkencave të Natyrës, 2: 89–93.
- Dhora Dh., 1978a. Gastropodë prozobrankë të rretheve Shkodër dhe Lezhë. Punim për fitimin e gradës shkencore “Kandidat i Shkencave”, 111 faqe.
- Dhora Dh., 1979b. Disa të dhëna mbi moluskun *Galba truncatula* (Müller 1774) në kushtet e vendit tonë. Buletini i Shkencave të Natyrës, 1: 91–96.
- Dhora Dh., 1981. Lloje të reja kërmijsh të rendit Stylomatophora të vendit tonë. Buletini i Shkencave të Natyrës, 2: 79–81.
- Dhora Dh., 1983. Mbi kërmillin *Viviparus mamillatus* Küster të vendit tonë. Buletini i Shkencave të Natyrës, 1: 101–103.

- Dhora Dh., 1985. Tre popullime të kërmillit *Cepaea vindobonensis* FER. (në Shkodër). Buletini i Shkencave të Natyrës 2: 105–108.
- Dhora Dh., 1988a. Të dhëna për dy lloje kërmijsh të nënfamiljes Helicigoninae. Buletini i Shkencave të Natyrës, 2: 95–97.
- Dhora Dh., 1988b. Të dhëna më të plota për dy kërmij të vendit tonë. Buletin Shkencor ILP, 2: 131–133. Shkodër.
- Dhora Dh., 1990. Kërmij dëmtues të kulturave bujqësore të rrethit të Shkodrës. Buletini i Shkencave të Natyrës 2: 95–97.
- Dhora Dh., 1994. Mbi popullatën e kërmillit *Theba pisana* (MULLER 1774) të Velipojës. Buletin Shkencor U Sh “Luigj Gurakuqi” 48, Seria e Shkencave Natyrore, fq. 45–48.
- Dhora Dh., 1995–1996. *Viviparus mamillatus* (KUSTER 1852) në pjesën shqiptare të Liqenit të Shkodrës. Buletin Shkencor U Sh “Luigj Gurakuqi” 48 (1), Seria e Shkencave Natyrore, fq. 47–52.
- Dhora Dh., 1995. List of molluscs of Shkodra Lake. Bios. (Macedonia, Greece) 3: 21–24.
- Dhora Dh., 1997–2000. Molusqe të tokës dhe ujërave të ëmbla të Shqipërisë. Buletin Shkencor U Sh “Luigj Gurakuqi” 49 (1), Seria e Shkencave Natyrore, fq. 47–61.
- Dhora Dh., 1997a. Molusqet e tokës dhe ujërave të ëmbël. Libri i Kuq (Bimë, Shoqërimi bimorë dhe Kafshë të rrezikuara), ShRMMNSh, ShRShGjSh, ShBSh, REC. fq. 121–134.
- Dhora Dh., 1997b. Molusqet. Kashta, L., Dhora, Dh. & Rakaj, M. Lista e bimëve dhe kafshëve të Liqenit të Shkodrës (pjesa shqiptare). Botim i Sektorit Shkencor të Bioekologjisë për Liqenin e Shkodrës, U Sh “Luigj Gurakuqi”, fq. 20–21.
- Dhora Dh., 1999a. Koregjime taksonomike dhe specie të kërcënuara të molusqeve të Liqenit të Shkodrës. Bio & Eko. Botim i Sektorit Shkencor të Bioekologjisë për Liqenin e Shkodrës, U Sh “Luigj Gurakuqi”, fq. 14–16.
- Dhora Dh., 1999b. Molusqet (endemike të Liqenit të Ohrit). Raporti Kombëtar “Strategjia dhe plani i veprimit për biodiversitetin”, Shtojcat / Anekset, C. Lista e taksoneve endemike, fq. 101.
- Diava Consulting, OIKON, Abkons, 2020. Final Report on the Study of Landscape of Llogara National Park in Albania, Enabling financial sustainability of Protected Areas System in Albania, project report.
- Fehér Z. & Drimmer L., 2004. Molluscs in the Megyeri Collection. Soosiana, 31: 21–25.
- Fehér Z., 2004. A revision of the genus *Cochlostoma*, subgenus *Titanopoma* (Gastropoda, Caenogastropoda, Cochlostomatidae), in particular the forms occurring in Albania. Basteria, 68: 25–44.
- Fehér Z. & Eröss Z., 2009. Contribution to the Mollusca fauna of Albania. Results of the field trips of the Hungarian Natural History Museum between 1992 and 2007. Schriften zur Malakozoologie, 25: 3–21.
- Fehér Z., Eröss Z. & Varga A., 2001. Contributions to the knowledge of the Albanian *Cochlostoma* fauna (Gastropoda: Cyclophoridae). Schriften zur Malakozoologie, 17: 71–78.
- Fehér Z., Eröss Z., Kontchán J. & Eröss Z., 2004. Collecting Sites Of The Zoological Expeditions Of The Hungarian Natural History Museum to Albania (1992-2003). Folia historico-naturalia Musei Matraensis, 28: 67–82.
- Goudeli G., Parmakelis A., Proios K., Anastasiou I., Radea C., Pafilis P. & Triantis K.A., 2021. The land snails of Lichadonisia islets (Greece). Ecologica Montenegrina, 39: 59–68.
- IUCN, 2011. IUCN Red List of Threatened Species (ver. 2011.1). Available at: <http://www.iucnredlist.org>. (Accessed: 30 June 2017).
- IUCN, 2020. The IUCN Red List of Threatened Species. Version 2020-3. Available at: www.iucnredlist.org. (Accessed: 10 December 2020).
- Jaekel S.H.F. & Schmidt H.A., 1961. Beitrag zur Molluskenfauna von Albanien. Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden, 26: 63–78.
- Jaekel S.H.F., Klemm W. & Meise W., 1957. Die Land- und Süßwasser-Mollusken der Nördlichen Balkanhalbinsel. Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden, 23: 141–205.
- Kümmerle J.B., 1926. Útívázlat - Reiseskizze. In: A Magyar Tudományos Akadémia Balkán-kutatásainak tudományos eredményei. (Teleki P. & Csiki E., Eds. 1926) Vol. 3. Adatok Albánia flórájához. Csiki Ernő, Jávorka Sándor és Kümmerle Jenő Béla gyűjtése - Additamenta ad floram Albaniae. Explorationes ab E. Csiki, A. Jávorka Et E. B. Kümmerle Peractae, The Hungarian Academy of Sciences, Budapest, pp. 8–89,
- Lika R., Deli T., Papparisto A., Eröss Z.P. & Feher Z., 2021. Contribution to the taxonomy and distribution of Aciculidae (Mollusca: Gastropoda: Caenogastropoda) in Albania and the surroundings. Zootaxa, 5004: 131–150. <https://doi.org/10.11646/zootaxa.5004.1.5>
- Martens E. Von, 1889. Griechische Mollusken. Gesammelt von Eberh Von Örtzen. Archiv für Naturgeschichte, 55: 169–240.
- Mousson A., 1859. Coquilles terrestres et fluviatiles, recueillies dans l’Orient par M. le Dr. Alex Schläfli. Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich, 4: 12–36, 253–297.
- Neubert E., Seddon M.B., Allen D.J., Arrébola J., Backeljau T., Balashov I., Bank R., Cameron R., de Frias Martins A.M., De Mattia W., Dedov I., Duda M.,

- Falkner G., Falkner M., Fehér Z., Gargominy O., Georgiev D., Giusti F., Gómez Moliner B.J., Groh K., Ibáñez M., Kappes H., Manganelli G., Martínez-Ortí A., Nardi G., Neiber M.T., Páll-Gergely B., Parmakelis A., Prié V., Reischütz A., Reischütz P.L., Rowson B., Rüetschi J., Slapnik R., Son M., Štamol V., Teixeira D., Triantis K., Vardinoyannis K., von Proschwitz T. & Walther F., 2019. European Red List of Terrestrial Molluscs: snails, slugs, and semi-slugs. IUCN: Cambridge, UK and Brussels, Belgium. <https://portals.iucn.org/library/node/48439>
- Riedel A. & Welter-Schultes F., 1996. Eine neue *Gyrulina*-Art aus Albanien (Gastropoda: Zonitidae). *Schriften zur Malakozoologie*, 9: 17–20.
- Sturany R. & Wagner A.J., 1915. Über schalentragende Landmollusken aus Albanien und Nachbargebieten. *Denkschriften der Kaiserlichen Akademie der Wissenschaften in Wien, Mathematisch-naturwissenschaftliche Klasse*. 91: 19–138, pl. 1–18, 1 Kart.
- Subai P. & Fehér Z., 2006. Revision of the Ariantinae, 3. *Superba* gen. nov., with the description of three new species (Gastropoda: Pulmonata: Helicidae). *Archiv für Molluskenkunde*, 135: 205–224.
- Subai P., 1997. Beitrag zur Kenntnis albanischer und montenegrinischer Ariantinae (Gastropoda: Helicidae). *Annales des Naturhistorischen Museum Wien* 99B: 521–534.
- Subai P., 1999a. Beschreibung einer neuen Art *Argna thracica*, sowie Angaben über die griechischen Argnidae. *Schriften für Malakozoologie* 13: 43–48.
- Subai P., 1999b. Beschreibung einer neuen Art der Gattung *Metafruticicola* (Gastropoda: Hygromiidae) aus NWGriechenland und Albanien. *Schriften für Malakozoologie*, 13: 49–54.
- Wagner A.J., 1913–15. Die Familia der Clausiliidae. In: Rossmässler's Iconographie der Land- und Süßwassermollusken mit vorzüglicher Berücksichtigung der europäischen noch nicht abgebildeten Arten. Neue Folge, (2) 21, pp. 1–65, C.W. Kreidel, Wiesbaden.
- Wagner A.J., 1919. Beschreibungen neuer oder bisher wenig gekannter clausiliiden (ii. teil) vorläufige Mitteilung über die von Dr. A. Penther in den Jahren 1914, 1916 und 1918 in Nordalbanien gesammelten Mollusken. *Anzeiger, Akademie der Wissenschaften in Wien, mathematisch-naturwissenschaftliche Klasse*, 56: 70–74.
- Welter-Schultes F.W., 2012. European non-marine molluscs, a guide for species identification: Bestimmungsbuch für europäische Land- und Süßwassermollusken. Planet Poster Editions, 674 pp.
- Welter-Schultes F.W., 1996. Non-marine molluscs recently collected in Albania. *Schriften zur Malakozoologie*, 9: 21–31.
- Westerlund C.A. & Blanc H., 1879. Aperçu sur le faune malacologique de la Grèce inclus l'Épire et la Thessalie. *Coquilles extramarines, Napoli*, 161 pp.
- Wohlberedt O., 1909. Zur Fauna Montenegros und Nordalbaniens (Mollusken, Käfer, Isopoden, Chilopoden, Diplopoden). *Wissenschaftliche Mitteilungen aus Bosnien und der Herzegowina*, 11: 585–722.